United States Department of Agriculture


Forest Service

Southwestern Region

MR-R3-16-3

Field Guide to Insects and Diseases of Arizona and New Mexico Forests







The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TTY). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, SW, Washington, DC 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TTY). USDA is an equal opportunity provider and employer.

Field Guide to Insects and Diseases of Arizona and New Mexico Forests

Mary Lou Fairweather, Plant Pathologist Joel McMillin, Entomologist Terry Rogers, Entomologist Dave Conklin, Plant Pathologist Bobbe Fitzgibbon, Entomologist

> USDA Forest Service Southwestern Region 333 Broadway Blvd., SE Albuquerque, NM 87102

www.fs.fed.us/r3/resources/health

2006

Acknowledgements

The authors thank all former pathologists and entomologists,

developed in the 1980s, or worked on getting this version started. Special thanks to Sandra Roberts, printing specialist and Dan Ryerson, GIS specialist for the Southwestern Region in Albuquerque, New Mexico.

Photographs and other illustrations were obtained, as

USDA Forest Service, and other sources as cited.

Preface

This book replaces "Forest Insect and Disease Field Guide" (Beatty 1983). Although amendments had been made to the former guide, our knowledge of forest insects and diseases has increased and

and/or economic impacts on forest resources such as recreation, wildlife, wood production, and watershed quality.

each with its own description, illustrations, and host index. Literature

Table of Contents

Roles of Forest Insects and Pathogens
Invasive Pests
Guide to Forest Insects and Diseases4
Forest Insect Defoliators
Western Spruce Budworm
11
Pine-feeding Needleminers
White Fir Needleminer
Pandora Moth18
Tiger Moth
New Mexico Fir Looper
Nepytia janetae
Fall Webworm
Western Tent Caterpillar
Large Aspen Tortrix
Alder Flea Beetle
Sap-Sucking Insects, Gall Formers, and Mites 37
Aphids
Cooley Spruce Gall Adelgid
Spruce Aphid
Pinyon Needle Scale
Prescott Scale
Pine Needle Scale
Ponderosa Pine Resin Midge50
Pinyon Spindle Gall Midge
Spittlebugs
Spider Mites54
Bark Beetles55
Pine Engravers
Western Pine Beetle
Mountain Pine Beetle
Roundheaded Pine Beetle

Southern Pine Beetle and Mexican Pine Beetle79	0
Red Turpentine Beetle	
Spruce Beetle	
Western Balsam Bark Beetle	
Fir Engraver	
Twig Beetles	
Cedar Bark Beetles	
Wood Borers8	
Wood Borers	
Juniper Borers	
Ambrosia Beetles9	2
Cone and Seed Insects9	4
Cone Beetles	
Ponderosa Pine Seedworm	
Pine Seed Chalcid	
Pine Coneworm	9
Bud and Shoot Insects	0
Bud and Shoot Insects	1
Bud and Shoot Insects	1
Bud and Shoot Insects10Pine Tip Moths10Western Pineshoot Borer10Pinyon Pitch Nodule Moth10	1 3 5
Bud and Shoot Insects10Pine Tip Moths10Western Pineshoot Borer10Pinyon Pitch Nodule Moth10Bark Moths10	1 3 5 6
Bud and Shoot Insects10Pine Tip Moths10Western Pineshoot Borer10Pinyon Pitch Nodule Moth10Bark Moths10Pitch Moth10	1 3 5 6
Bud and Shoot Insects10Pine Tip Moths10Western Pineshoot Borer10Pinyon Pitch Nodule Moth10Bark Moths10	1 3 5 6
Bud and Shoot Insects 10 Pine Tip Moths 10 Western Pineshoot Borer 10 Pinyon Pitch Nodule Moth 10 Bark Moths 10 Pitch Moth 10 Juniper Twig Pruner 10 Insects of Wood Products 10	1 3 5 6 6 7 8
Bud and Shoot Insects 10 Pine Tip Moths 10 Western Pineshoot Borer 10 Pinyon Pitch Nodule Moth 10 Bark Moths 10 Pitch Moth 10 Juniper Twig Pruner 10 Insects of Wood Products 10 Carpenter Ants 10	1 3 5 6 6 7 8 9
Bud and Shoot Insects 10 Pine Tip Moths 10 Western Pineshoot Borer 10 Pinyon Pitch Nodule Moth 10 Bark Moths 10 Pitch Moth 10 Juniper Twig Pruner 10 Insects of Wood Products 10	1 3 5 6 6 7 8 9
Bud and Shoot Insects 10 Pine Tip Moths 10 Western Pineshoot Borer 10 Pinyon Pitch Nodule Moth 10 Bark Moths 10 Pitch Moth 10 Juniper Twig Pruner 10 Insects of Wood Products 10 Carpenter Ants 10	1 3 5 6 6 7 8 9
Bud and Shoot Insects 10 Pine Tip Moths 10 Western Pineshoot Borer 10 Pinyon Pitch Nodule Moth 10 Bark Moths 10 Pitch Moth 10 Juniper Twig Pruner 10 Insects of Wood Products 10 Carpenter Ants 10 Termites 11	1 3 5 6 6 7 8 9 1
Bud and Shoot Insects 10 Pine Tip Moths 10 Western Pineshoot Borer 10 Pinyon Pitch Nodule Moth 10 Bark Moths 10 Pitch Moth 10 Juniper Twig Pruner 10 Insects of Wood Products 10 Carpenter Ants 10 Termites 11 Foliar Diseases 11 Hardwood Leaf Diseases 11 Black Leaf Spot 11	1 3 5 6 6 7 8 9 1 3 4
Bud and Shoot Insects 10 Pine Tip Moths 10 Western Pineshoot Borer 10 Pinyon Pitch Nodule Moth 10 Bark Moths 10 Pitch Moth 10 Juniper Twig Pruner 10 Insects of Wood Products 10 Carpenter Ants 10 Termites 11 Hardwood Leaf Diseases 11 Black Leaf Spot 11 Ink Spot Leaf Blight 11	1 3 5 6 6 7 8 9 1 3 4 5
Bud and Shoot Insects 10 Pine Tip Moths 10 Western Pineshoot Borer 10 Pinyon Pitch Nodule Moth 10 Bark Moths 10 Pitch Moth 10 Juniper Twig Pruner 10 Insects of Wood Products 10 Carpenter Ants 10 Termites 11 Foliar Diseases 11 Hardwood Leaf Diseases 11 Black Leaf Spot 11 Ink Spot Leaf Blight 11 Melampsora Rust 11	1 3 5 6 6 7 8 9 1 3 4 5 6
Bud and Shoot Insects 10 Pine Tip Moths 10 Western Pineshoot Borer 10 Pinyon Pitch Nodule Moth 10 Bark Moths 10 Pitch Moth 10 Juniper Twig Pruner 10 Insects of Wood Products 10 Carpenter Ants 10 Termites 11 Hardwood Leaf Diseases 11 Black Leaf Spot 11 Ink Spot Leaf Blight 11	1 3 5 6 6 7 8 9 1 3 4 5 6 7

Conifer Needle Diseases	20
Lophodermella Needle Cast	21
Elytroderma Needle Cast	23
Red Band Needle Blight	25
Rhabdocline Needle Cast	26
True Fir Needle Cast	28
White Pine Needle Cast	29
Pinyon Needle Rust	30
Stem Decays and Stains	32
Red Ring Rot	
Red Rot	36
Indian Paint Fungus	39
Red Belt Fungus	41
Pouch Fungus	42
Blue Stain Fungi 1	43
False Tinder Conk	44
Heart Rots of Oak	45
Canker Rot of Oak	46
Cankers 1	47
Black Canker	48
Sooty Bark Canker	50
Cytospora Canker	52
Cryptosphaeria Canker	54
Hypoxylon Canker	56
Atropellis Canker	58
Stem and Cone Rusts of Pine	60
White Pine Blister Rust	61
Limb Rust1	64
Western Gall Rust	66
Comandra Blister Rust	67
Gymnosporangium Rusts	69
Fir Broom Rust	71
Spruce Broom Rust	71
Southwestern Cone Rust	75

Mistletoes
Dwarf Mistletoes
True Mistletoes
Root Disease
Armillaria or Shoestring Root Rot
Annosus Root Rot
Tomentosus Root Disease
Schweinitzii Root and Butt Rot
Black Stain Root Disease
Ganoderma Root Rot
Non-infectious Disorders
Drought
Frost
Winter Injury
Red Belt
Windthrow
Lightning
Salt Damage
Burls
Animal Damage
Ungulates
Black Bear
Porcupine
Sapsucker
Glossary
Photography and Illustration Credits
Instructions for Submitting Insect and Disease
Selected References
Host Index
General Index

Roles of Forest Insects and Pathogens

Most forest insects and pathogens in the Southwest are naturally occurring components of ecosystems and play an important role in dynamic processes. They affect short- and long-term vegetative structural diversity, provide food and habitat for animals, and contribute to biological diversity. These organisms, along with

Outbreaks of forest insects and diseases can result in shifts in forest composition and structure. The degree of shift depends upon the particular insect and/or disease and on the condition of the ecosystem affected.

The actions of insects are often easily observed as their populations can quickly expand under favorable climatic conditions.

altering forest succession. For example, bark beetle populations can increase dramatically under drought conditions and cause large-scale tree mortality. In contrast, indigenous tree pathogens increase gradually and their presence and effects may go relatively unnoticed. Although most people see dwarf mistletoe in our forests, few realize that it is the primary agent of growth loss and will gradually kill trees. Insects and pathogens often interact, as is seen when bark beetles target diseased trees.

A key to insect and disease damage is provided in the "Guide to Forest Insects and Diseases" section, directing the reader to the appropriate insect or disease section. There is also a host index in the back of the book. A description for each insect and disease includes hosts, damage, symptoms, biology, and effects or impacts. References are also provided.

Please refer to your local Forest Service, Forest Health

measures.

References: 16, 50, 58, 98, 99, 105

Invasive Pests

Similar to the rest of the Nation, Arizona and New Mexico's forests and grasslands are threatened by invasions of exotic species of insects, disease pathogens, and plants. Over the past 200 years, several thousand foreign plant and animal species have been introduced in the United States. About one in seven of these species has become invasive.

Two invasive pest species that were introduced into the United States in the early 1900s have become established in forested lands of New Mexico and Arizona. White pine blister rust, caused by *Cronartium ribicola*, was discovered in southern New Mexico in 1990, and is now found throughout most of the range of its host in the Sacramento and adjoining White Mountains. Infected




Figure 1. Adult spruce aphid

white pines have also been found in nearby Capitan Mountains and Gallinas Peak, New Mexico. In 2005, a population was observed on the Gila National Forest, New Mexico, about 3 miles from the

became established around 1970 on the west side of the Sacramento Mountains, but it is not clear how the fungus was introduced. Over time, this disease will have a major impact on white pines in the Sacramento Mountains and some other parts of the Southwest.

The second species introduced in the region is spruce aphid, *Elatobium abietinum* ornamental tree in the city of Santa Fe, New Mexico, and in a natural forest setting in the late 1980s. Its distribution extends from the San Francisco Peaks in northern Arizona, to the Sacramento Mountains of southern New Mexico. The damage caused by this insect in the Southwest is much greater than that observed elsewhere in the

United States, which may be due to variations in the insects' behavior. Research in the Southwest found that spruce aphid populations increase in the fall, have a sexual life cycle, and a greater cold-hardiness. all factors that



Figure 2. White pine blister rust on ribes

may be contributing to the insect's success.

The Animal and Plant Health Inspection Service (APHIS) is responsible for regulating the movement of plants and plant materials that may carry pest organisms, and for detection and eradication of new pest introductions. The USDA Forest Service, the Bureau of Indian Affairs, and other Federal land managing agencies are responsible for reducing the impact of invasive exotic plants on lands they manage. Despite the efforts, we expect increased introductions of nonnative species as well as accidental introductions of native species of the U.S. (from one part of the country to another) due to expansion of world and national trade. Early detection and rapid response are important to minimizing invasive species establishment and subsequent resource damage.

References: 13, 56, 105

Guide to Forest Insects and Diseases¹

Damage confined to leaves, buds and shoots:

Defoliating Insects. Affected needles with a reddish cast,
insects or signs of their feeding present. Needles and buds are
partially or totally consumed. Caterpillar or beetle-like insects
usually present on foliage in early spring or summer. Leaves chewed,
mined, skeletonized or webbed. Viewed from a distance, affected
trees have a brownish or yellowish cast7
Sap-Sucking Insects and Mites. Curled or discolored
foliage. Whole insects or cast skins and small (1-2 mm long).
Cottony tufts, galls, and honeydew sometimes present37
Galls Caused by Insects. Abnormal, misshapen, swollen or
spherical growth on needles or leaves, or in the bark or twigs. Insect
larvae or frass present inside the galls52
Bud and Shoot Insects. Malformed, enlarged buds on
young growth, commonly encountered on terminal buds of branches.
In some cases, multiple leaders result100
Foliar Diseases (including needles). Foliage yellow, brown,
or spotted (insects largely absent); orange, yellow, or black pustules
or other fungal fruiting bodies present. Fruiting bodies are very
small113
Abiotic Disorders. Foliage or buds yellowing, browning, or
dying after exposure to early or late frosts, air pollutants, drought,
or winter damage; entire crown may show signs of stress or die as a
unit; cause of the damage may not be apparent but signs of insect or
disease agents are generally absent204

 $^{^{\}rm I}$ Adapted from "Insects and Diseases of Alaskan Forests" (Holsten et al., 2001)

spherical growth in the cones or seeds. Insect larvae or frass present
inside the galls99 Moths and Midges. External evidence of insect activity is
found within the cone axis, cone scales, and seeds. Boring dust is
Cone Rust. Enlarged cones on Chihuahua pine, with orange spores under papery outer covering175
Damage confined to branches and stems:
Witches' Brooms.
branches infected with mistletoe or fungi.
Needles retained on those caused by dwarf mistletoe177
Needles cast on those caused by fungi, only to reappear the
next year123
Cankers and/or Animal Damage. Large patches of dead
bark on live trees, sometimes exposing wood; often on main tree
bole; fruiting bodies of fungi may be present147
Strips or sections of bark removed by animals, fungi follow
wounding213
Stem Decay. Wood is hollow, pitted, spongy, breaks into
crumbly pieces, or is stained in color. Fruiting bodies of fungi may
be present. Older trees, or those physically injured in the past usually
affected132
Bark Beetles. Insect activity characterized by boring dust

and/or resin on bark and around base of trees. Galleries commonly found under bark and usually less than 0.6 cm in width, at times into the sapwood, rarely penetrating the wood. Boring dust reddish to

Galls Caused by Insects. Abnormal, misshapen, swollen, or

Wood Borers. Larval borings are usually granular. Galleries
penetrate into the sapwood and, at times, into heartwood of
weakened or recently killed trees or logs87
Abiotic Disorders. Globe-shaped, large, woody growths or
swellings on branches or stems of ponderosa and various hardwoods
(burls)212

Damage confined to roots and lower stem:

Noninfectious Disorders. Dead and dying trees found in small or large groups, not usually solitary trees. Entire crown of tree shows signs of stress and dies as a unit similar to damage by root disease. Fruiting bodies of pathogenic fungi usually absent, but weakly pathogenic and saprophytic fungi (those restricted to dead tissues) and secondary insects may be present. ------204

efoliating insects damage trees by eating leaves or needles, removing the photosynthetic tissue critical

of leaves or needles results in growth loss, increased susceptibility to attack by other insects and pathogens, and sometimes tree mortality.

The impact of defoliation on individual trees is dependent on a variety of factors. Tree species vary in their tolerance to defoliation. In general, hardwood species can sustain repeated defoliation over several years because they store large food supplies and can refoliate in the same year. The timing of the defoliation relative to seasonal growth phenology is also important. For example, late season defoliation of hardwoods has a lower impact than does late season defoliation of conifers. A single late season defoliation of pines often results in tree mortality. Trees that are healthy and growing vigorously will generally survive defoliation better than stressed trees.

Historically, the two most destructive defoliators in the Southwestern Region were the western spruce budworm and the

growth loss, top-kill, increase susceptibility to bark beetles, and cause outright tree mortality. More recently, three other defoliators have been causing heavy defoliation in higher elevation forests. Beginning in the early 1990s, the exotic spruce aphid and a native looper, *Nepytia janetae*, are causing high levels of mortality in the

Sacramento Mountains of southeastern New Mexico causing elevated

Many other defoliating insects that have the potential to cause localized problems are not included in this guide. Consult Federal or State forest health personnel for assistance on insects not described in this guide.

References: 40, 66, 72, 108

Western Spruce Budworm Choristoneura occidentalis (Freeman)

Hosts:

Symptoms/signs: Adults are a mottled orange brown and have a wing span of 22 to 28 mm. Eggs are white to light green in color and are laid in shingle-like masses on the underside of needles. Newly hatched larvae are cream colored with brown heads. Full-grown larvae are 25 to 32 mm long with brownish head and body

and prominent ivory colored spots. Pupae are 12 to 16 mm long, broad at the head and tapering toward the tail. Signs of feeding include current foliage being partially or fully chewed over the entire tree; expanding buds mined and evidence of feeding on second and third year shoots possible;



Figure 3. Adult western spruce budworm with pupal case.

Figure 4. Egg mass of western spruce budworm on underside of needle. Note shingle-like feature.

and shoots webbed into feeding shelters giving the tree a reddish-brown appearance.

Biology: Eggs are laid in July and August and hatch in about 10 days. The initial larvae do not feed, but spin silken shelters (hibernacula) under

bark scales or lichens where they will hibernate. The next spring larvae mine old needles until the buds swell and then bore into the buds and feed on the expanding needles. Later they web the growing tips together and feed on the new needles. The larvae pupate in June and July. Adults begin to appear in July and start egg laying for the next generation.




Figure 5. Late instar larva of western spruce budworm feeding on foliage.

Effects: Defoliation

by western spruce budworm can cause growth loss after 1 or 2 years. Repeated heavy defoliation (4 or 5 years) can cause a




Figure 6. Defoliation of Douglasfir tree caused by western spruce budworm.

tree deformity, top-killing, and ultimately tree mortality, particularly in seedlings and saplings.

Similar Insects and Diseases:

needleminer, *Epinotia meritana*, feeds inside the needles of true

yellowish color and do not have a chewed appearance. The mature larvae are much smaller (8mm) than western spruce budworm and yellowish green to cream colored. The adult

moths have wings that are dusty gray with alternating bands of dark needleminer can cause branch mortality, deterioration of tree crowns




Figure 7. Near view of western spruce budworm defoliation to Douglas-fir.

Douglas-fir Tussock Moth Orgyia pseudotsugae (McDunnough)

Hosts: Symptoms/signs:

moth caterpillar is grayish with brightly colored tufts of hair and a shiny black head. There are also two long horns of black hairs behind the head and another at the rear of the body. Insect feeding results in needle damage and defoliation, which is sometimes followed by





Figure 8. Adult male (left) and female (right) Douglas-fir tussock moth.

top-kill. Small cocoons and egg mass cases are attached to the underside of twigs and branches. Biology: Eggs hatch from mid-May to early June and caterpillars feed on the current year's developing foliage. Figure 9. Adult Young female Douglaslarvae are fir tussock moth 4 to 7 mm on egg mass. Note long and that female moth covered is wingless. with long,

Figure 10. Egg mass and early instar larvae of Douglas-fir tussock moth.



thin body hairs that later develop into tufts. Because the female moth is wingless, the primary means of dispersal from tree to tree is by windblown larvae. Young larvae congregate on the tops of defoliated trees and drop by silken threads that may be over 3 meters long. These threads eventually break from the tree and give a ballooning effect to the larvae. Mature larvae are about 20 to 30 mm long with a gray or brown body and a shiny black head. Two long hair pencils project forward from behind the head and another occurs at the rear of the body. Four brushes of brown or cream-colored hairs

Some people develop an itchy rash from exposure to the frequently airborne caterpillar hairs. After the caterpillars are about halfgrown, they feed on all age classes of needles throughout the crown. Pupation occurs inside a thin, silk cocoon spun on the undersides of branches from late July to the end of August. Adults emerge shortly

and branches.

Effects:

both overstory and understory trees relatively quickly (1 or 2 years). Top-kill occurs in less severe outbreaks. Bark beetle epidemics can occur in tussock moth defoliated areas.

Similar Insects and Diseases: Early damage and webbing in trees may be similar to that caused by western spruce budworm; however, the larvae are very distinct.

References: 9, 23, 111

Pine-feeding Needleminers Coleotechnites spp.

Hosts:

Symptoms/Signs: Adults of pine-feeding needleminers are small, narrow-winged, mottled, silvery-gray moths with a wingspan of about 10 mm. Eggs are too small, 0.2 mm in diameter, to be noticeable. Larvae are brown, 8 mm long when fully developed, and usually found mining within current year's needles. The pupae

are elongate, cylindrical black, and about 6 mm long. Needles inhabited by third and fourth instar larvae turn a faded yellowbrown in color and have several tiny holes in them for frass disposal and larval exit.




Figure 14. Silhouette of Coleotechnites larvae feeding inside ponderosa pine needle.

Biology:

are laid inside previously mined needles. Larvae begin hatching in July, crawl to and mine into previously uninfested needles and overwinter. Pupation occurs in late spring.

Effects: Persistent infestations can cause severe discoloration, defoliation, and reduced growth of stems, shoots, and needles.



White Fir Needleminer Epinotia meritana (Heinrich)

Host:

Symptoms/Signs: Larvae mine the needles, resulting in bleached-yellow mined needles from late spring to early fall. Mature larvae are about 8 mm long, yellowish-green to cream colored, with brown to black heads. Pupae are orange to dark brown, about 5.5 mm in length, and can often be seen protruding from the hole in the mined needle prior to emergence. The adults are 11 mm dusty gray moths. They can be seen swarming around host trees in late June and July.

Biology:

per year in New Mexico and Arizona. Eggs are deposited, usually one egg per needle, in late June and July. Eggs hatch in August and September. Young larvae immediately bore into the needle where they overwinter. As weather warms in the spring, feeding begins.

three-year-old needles are also mined during outbreak conditions.

Figure 17. Defoliated white fir caused by the white fir needleminer.




Figure 18. Closeup of white fir needleminer defoliation to white fir.

Two to six needles are usually webbed together and remain on the tree for the duration of the summer. Pupation occurs within the last mined needle in June or early July. Adult moths emerge in 10 to 14 days after pupation. Mating occurs within 2 to 3 days of emergence.

Effects: After several consecutive years of heavy defoliation a high proportion of limb mortality and widespread tree mortality may occur. This can give individual trees or even entire stands a silvery appearance. These heavily defoliated stands decline in vigor

Similar Insects and Diseases: Similar defoliation may be

References: 23, 109

Pandora Moth Coloradia pandora (Blake)

Hosts: Ponderosa pine
Symptoms/Signs: Adult
pandora moths are very large and
heavy bodied, about 2.5 to 4.0 cm
long, with a wingspread of 7 to 11
cm. The forewings are brownish
gray and hindwings are light pinkish
gray, each marked with a black dot
and a dark wavy line. The males are
distinguished by having large, feathery antennae. The globular eggs,
bluish green to bluish gray, are deposited in clusters of 2 to 50. Early
instar larvae are about 5 mm long. They have shiny black heads and
black to brownish bodies that are covered with short, dark hairs.

Fifth instar larvae grow to about 6 to 8 cm long and are brown to yellowish

the ground where they pupate. They remain in the pupal stage for the next 12 to 13 months.

Effects: During outbreaks, defoliation can be severe over large areas. Due to the 2-year life cycle, however, defoliation occurs only during alternate years. Outbreaks on the Kaibab Plateau have subsided without causing any lasting damage. However,



Figure 22. Pupae of pandora moth.

some growth loss and even mortality can occur especially if trees are severely stressed from additional factors such as drought or heavy dwarf mistletoe infections.

Similar Insects and Diseases: May be confused with




Figure 23. Defoliation of ponderosa pine caused by pandora moth on the Kaibab NF, Arizona.

bodies, are smaller, 18 to 25 mm long, and have eight pairs of leglike appendages on the abdomen. Defoliation

usually occurs on an individual or small group of trees and is not widespread like that of a pandora moth outbreak.

References:

10, 23, 85

Pine Butterfly Neophasia menapia (Felder and Felder)

Host: Ponderosa pine **Symptoms/signs:** Newly

gregarious feeders on older foliage and then move to the new foliage as they mature. Mature larvae are approximately 2.5 cm long, dark green in color and have light green heads. The body is covered with

stripes on the side. Adult moths are white with black markings.

Biology:

and lay their green eggs from August through October. The eggs overwinter and the larvae hatch the following June or about the time new foliage appears. The small larvae feed in clusters with




Figure 24. Adult pine butterfly.

their black heads oriented toward the tip of the needle. As the larvae mature they feed singly. They pupate in late July for approximately 15-20 days.

Figure 25. Larva of pine butterfly.

Effects: This insect has caused spectacular defoliation in

it causes little defoliation in the Southwest.

Similar Insects and Diseases:

resembles the common cabbage

larvae feeding in groups may be confused with other gregariously

These would include *Neodiprion* spp. and *Zadiprion* spp. of is also a defoliator of ponderosa pine. This insect has a 2-year life cycle. Mature larvae are distinctly larger, brown to yellowish green in color and possess stout, branched spines on each body segment.

References: 23

Figure 26. Pupa of pine butterfly.

Pine Sawflies Neodiprion spp., Zadiprion spp.

Hosts:

Symptoms/signs: Pine

by species and by larval instar, but most are green or yellowish green in color with black, tan or orange head capsules. Larvae are found in either spring-summer or fall-winter feeding gregariously on




Figure 28. Row type egg mass of pine sawflies on ponderosa pine.

feed singly and consume most of the needle. Eggs are laid in slits cut in the edge of living pine needles. A papery cocoon covers pupae. Adults are broad waist wasps. Infested trees have sparse foliage and thin crowns.

Effects:

are found on ponderosa pine. Different species have different preferences for the size of host attacked, and location on the

Southwest typically attack open-grown trees




Figure 27. Adult female pine sawfly (Neodiprion spp.) ovipositing on ponderosa pine.

older foliage, consuming only the outer needle tissue while leaving the central ribs intact. The central ribs later turn yellow brown and break

off. Later instar larvae




Figure 29. Late instar of Neodiprion gillettei feeding on ponderosa pine.

or areas where pine is growing at a low density. The same trees are frequently defoliated year after year. In general, defoliation causes slower growth. Repeated defoliation can result in top-kill and tree mortality.

Similar Insects and
Diseases: See pandora moth and

References: 18, 23, 63, 65, 67, 68, 69

Figure 30. Late instar larva of Zadiprion rohweri feeding on piñon pine.

Figure 31. Late instar larvae of Zadiprion townsendii feeding on ponderosa pine.

Note: The taxonomy of this group has recently changed. The *Neodiprion fulviceps* Complex, described in Furniss

Figure 32. Defoliated ponderosa pine caused by pine sawflies near Flagstaff, Arizona.

and Carolin, has been split into *N. fulviceps* and a new species *N. autumnalis*. The life cycle information described for *N. fulviceps* in Furniss and Carolin is now considered that for *N. autumnalis*.

Tiger Moth Lophocampa ingens (Edwards) (=Halisidota ingens)

Hosts:

Symptoms/Signs: Larvae feed on foliage and make large silken webs (tents) in the upper branches of host pines. Mature caterpillars are about 4 cm long, reddish brown to black in color and have tufts of black and yellow hairs on their back. Adult moths have dark forewings with large, white splotches and white hindwings.

Figure 33. Adult tiger moth.

Biology: The tiger moth has one generation per year in the Southwest. Adult moths emerge and lay eggs in August. During September and October caterpillars hatch from eggs, begin feeding on pine needles and producing webbing. Larvae overwinter in the webs in groups. In April and May, larvae resume feeding and expand their webs. At this point the insects and the webs become very noticeable. In June pupation occurs.

Tiger moths are also found feeding and forming tents on

Figure 34. Larvae of tiger moth.

It is believed that populations of this insect usually remain at low levels due to the action of predators, parasites, diseases, and cold winters.

Effects: Larvae feed gregariously in webs primarily

pine. Although the webs and larvae can be very noticeable in spring, this insect causes minor defoliation. Permanent tree injury rarely results from feeding that is usually limited to the upper foliage.

Similar Insects and Diseases: Tents are similar to those formed by tent caterpillars such as the western tent caterpillar; however, *L. ingens* is the most common moth to make




Figure 35. Tents and damage of ponderosa pine caused by tiger moth.

and white pines. Another species of tiger moth, *L. argentata subalpina*Rocky Mountain region.

References: 23, 104

New Mexico Fir Looper Galenara consimilus (Heinrich)

Hosts:

Symptoms/Signs: Larvae feed on the foliage often stripping its host of needles. The immature stages of this insect have not been described. The forewings of the adults are light gray overlain with grayish-brown and brownish-black scales.

Biology: Little is known about the biology of this insect. Outbreaks of this occasional pest occurred in 1924, 1928, 1951, 1958,



Figure 36. Larva of New Mexico fir looper.

1974, 1977, and 2002 in New Mexico, the only state in which this insect is known to occur.

Effects: This insect can cause heavy defoliation of Douglas-

heavy defoliation can result in high levels of tree mortality.





Figure 37. Damage of fir caused by New Mexico fir looper.

Figure 38. Near view of damage of fir caused by New Mexico fir looper.



Nepytia janetae (Rindge)

Symptoms/signs: Larvae of this insect feed on foliage, chewing needles of all ages. Defoliation can be light to heavy. Partially consumed needles can be found beneath the trees. Mature caterpillars are approximately 2.5 cm in length. The

upper surface of the larva is gray with

Hosts: Engelmann spruce,



Figure 39. Adult Nepytia janetae.

and green markings in a herringbone pattern and with irregular pale cream lateral stripes. The head capsule is mottled with a distinct black stripe above the mouthparts.

dark brown

Figure 40. Larva (looper) of Nepytia janetae.

These larvae

can be seen in the foliage, hanging from silken threads, beneath the tree on ground or snow and climbing up the bole of the tree. Adults

green pupae or empty pupal cases can be seen on the foliage wrapped in webbing. Mummies of parasitized larvae can be seen in the foliage attached to needles.

Biology: *N. janetae* has one generation per year. Adults appear in late June,

early July. Eggs hatch in late September. Larvae feed throughout the winter and spring. Some are present in larval form until early July. There is considerable overlapping of life stages.

Effects: During




Figure 41. Defoliated fir caused by Nepytia janetae in the White Mountains, Arizona.

be heavily defoliated resulting

in growth loss and, with multiple years of defoliation, tree mortality.




Figure 42. Near view of fir defoliated by Nepytia janetae in the White Mountains, Arizona.

Mortality can be due to defoliation alone or due to secondary infestation by spruce beetle and western balsam bark beetle. This insect caused extensive damage and

of Arizona in the late 1990s. Associated spruce beetle mortality occurred only in the

Similar Insects and Diseases:

Although there are similar insects in other regions (e.g. western hemlock looper and western false hemlock looper), none have been observed in the Southwest.

References: 56, 80

Fall Webworm Hyphantria cunea (Drury)

Hosts: Willow, alder, ash, chokeberry, cottonwood and madrone

Symptoms/signs:

Larvae feed on foliage, forming large webs in the branches of trees. Webs are noticeable in the fall when larval feeding takes place. There are two races: a blackheaded or northern race and a redheaded or southern race.




Figure 43. Adult and pupa of fall webworm.

Larvae of the southern race are

yellowish-tan with red or orange colored heads and brownish hair that arises from reddish-brown tubercles. Larvae of the northern race have a black head with a pale yellowish or greenish body that has a dark stripe on the back, and long white hairs rising from red




Figure 44. Larva of fall webworm.

or black tubercles. The black-headed race predominates in the West. Adults are white in color with orange markings on the body and legs. The wings have some black spots and a wing expanse of approximately 30 mm.

Biology:

Fall webworm has

one generation per year. Adults appear and lay eggs in late June and early July. The eggs hatch and the small larvae feed on both leaf

surfaces while larger larvae will consume the whole leaf. Larval feeding continues until mid-September. The insect overwinters in the pupal stage in a transparent cocoon in the soil, leaf litter, or on tree trunks.

Effects: This insect causes minor defoliation in most forested situations. It can cause loss of visual quality in ornamental plantings.

Similar Insects and Diseases: The western tent caterpillar is sometimes confused with the fall webworm, due to both having dark heads and dark stripes down their backs. However, western tent caterpillars feed in the spring while fall webworm feeds in the fall.




Figure 45. Damage caused by fall webworm.

Western Tent Caterpillar Malacosoma californicum (Packard)

Hosts: Aspen, willows, cottonwoods, and mountain mahogany

Symptoms/signs:

Western tent caterpillar is an early season defoliator with feeding damages typically occurring between May and June. Symptoms include moderate to complete defoliation of trees; large silken tents on branches:



Figure 46. Adult western tent caterpillar and egg mass.

and presence of larvae in and around the tents. Trees repeatedly defoliated will have sparse foliage, minor branch dieback, and in

some cases, tree mortality.

Mature larvae are 4 to 5 cm long and vary widely in coloration. Their heads are blue to black and body color patterns are mixtures of black, orange, and blue. Larvae are usually quite hairy.

Biology: Larvae



Figure 47. Larva of western tent caterpillar.

inside egg masses glued around twigs. Larvae emerge

from egg masses in spring and construct silken tents on branches that are used for shelter and molting during the daytime. At night, caterpillars feed outside of the tents. As the larvae mature, they disperse and become solitary feeders. Moths emerge from cocoons and glue egg masses to live twigs that are less than 2 cm in diameter.

Effects: Heavy defoliation of aspen for a number of years will cause growth loss and branch dieback. Some mortality may also occur during prolonged outbreaks. Outbreaks, however, are generally short lived, generally lasting 2 to 3 years.

Similar Insects and Diseases: See fall webworm and foliar diseases.

References: 3, 15, 23

Figure 48. Tents of western tent caterpillar on aspen.

Figure 49. Western tent caterpillar larvae and defoliation of aspen on the Carson NF, New Mexico.

Large Aspen Tortrix Choristoneura conflictana (Walker)

Host: Aspen

Symptoms/Signs: Presence of

foliage, partially eaten rolled-up leaves in which small, green to black larvae are feeding or pupating, incomplete defoliation before the buds have expanded in the spring.

Biology: Pale green egg masses

surfaces of the foliage in June and July. Second-instar larvae overwinter in silken hibernacula in bark crevices and other




Figure 50. Adult large aspen tortrix.

out-of-the-way places. The small larvae begin to migrate up the stems of aspen on warm days in early spring. The larvae mine the buds, sometimes causing complete defoliation before the buds open. Later stage larvae roll the leaves into shelters within which they feed and eventually pupate. The moths have a wingspread of 25 to 30 mm. The forewing

Figure 51. Closeup of damage caused by large aspen tortrix and larva.

is grayish with basal, middle, and outer brownish patches. There is one generation annually.

Effects: During outbreaks, this insect can completely defoliate aspen stands. Outbreaks characteristically last 2 to 3 years and can result in growth loss and some twig and branch dieback.

Similar Insects: Larvae and pupae are similar to those of the western spruce budworm. However, the larvae of the large aspen tortrix are green to grayish-black with a black head capsule while the western spruce budworm larvae are white to grayish-tan with a tan




Figure 52. Large aspen tortrix defoliation of aspen in Colorado.

Alder Flea Beetle Altica ambiens (LeConte)

Hosts: Alder
Symptoms/signs: Larvae
are skeletonizers, meaning that
they feed on the tissue between
the veins, but leave the veins
intact. Adults chew holes in the
leaves. Adult beetles are dark
shiny blue, and about 5 mm long.
The mature larvae are a little

longer and narrower than the adults, brown to black above and

vellowish below.



Figure 53. Adult alder flea beetle.

Biology: One generation is reported per year. Adults hibernate during the winter in duff at the base of trees and in other sheltered places. They emerge in early spring to resume feeding.




Figure 54. Early instar larvae of alder flea beetle on underneath side of alder leaf.

Eggs are laid in clusters on foliage. Larvae hatch and begin feeding within a few days. One source reports that larvae mature in August; but in the Southwest they mature earlier in the season.

Effects: No longterm effects are documented. Outbreaks are generally short lived and sporadic with heavy defoliation. Trees tend to recover quickly.

Similar Insects and Diseases: Although other defoliators feed on alder, outbreaks of these other insects have not been reported.




Figure 56. Defoliation of alder caused by alder flea beetle in Arizona.

he majority of sap-sucking insects are in the orders Hemiptera (true bugs) and Homoptera (aphids, leaf and plant hoppers, and scales). Most of these insects are relatively small in size and injure the host in two ways: (1) directly by sucking the host of part of its food supply and water, producing necrotic spots in host tissue, and (2) indirectly by introducing plant diseases.

The mouthparts of these insects and mites are formed into beak-like structures that are used to pierce host tissues and suck the sap. Damage by sap-sucking insects is often mistaken as a pathogen induced disease. A few of the sap-sucking insects are able to kill their hosts outright, but most reduce growth rates and weaken the tree. Trees injured by these insects may succumb to secondary insects or fungal diseases. Signs of sap-sucking insect injury consist of enlarged growths or galls, leaf curling, bleaching, or yellowing of foliage. Conifers are more severely injured than hardwoods.

In the Southwest, there are several sap-sucking insects that cause noticeable damage to trees and shrubs, but they have not affected forests at the landscape scale. Their impacts are typically more important to shade trees and ornamentals. However, a spruce aphid, *Elatobium abietinum*, has been severely impacting spruce forests and ornamentals in some areas of the Southwest since the 1990s.

as these organisms are quite small.

References: 23, 40

Aphids Aphididae, Adelgedae

Hosts: Hardwoods and conifers **Symptoms/signs:** The best evidence of infestation is the insects themselves or shiny leaves from the honeydew. Honeydew is a clear, sugary, sticky liquid that aphids excrete as they feed. It accumulates on the foliage, twigs, trunk and ground. Honeydew attracts ants, and often the presence of ants on a tree may indicate that aphids are there. It is also a good growth medium for sooty mold, which grow on affected portions of the tree. Aphids are small soft-bodied gregarious insects. Adults may be winged or wingless. Nymphs are wingless. They vary in color by species, but may be nearly colorless to green to black.




Figure 57. Cinara aphids on pine host.

Figure 58.
Eggs of
aphids on
pine needle.

Biology: Typically aphids produce several generations per year, most are parthenogenetic. The last generation is usually sexual, and then winter is spent in the egg stage. Different aphids feed on different parts of their hosts, including the foliage, twigs, trunk, and roots. All aphids have piercing mouthparts that they use to feed on sap from their host plants.

Effects: Feeding can result in formation of small necrotic spots. Heavy feeding by some species can cause early leaf drop, yellowing of foliage, and reduced growth. In general, these insects are not considered economically important in forest situations. In urban areas, the honeydew can accumulate on walkways and vehicles and attract the attention of the public.

Similar Insects and Diseases: Damage caused by aphids may resemble that caused by needle midges, other sucking insects, or needle diseases.




Figure 59. Aphid damage to conifer host.

Cooley Spruce Gall Adelgid Adelges cooleyi (Gillette)

Hosts:

Symptoms/signs:

Cone-like galls, 12-75 mm long, are formed on spruce branch tips as a result of feeding by the nymph stage of this insect. In late spring, galls are light green to purplish in color and the nymphs may be found inside. In late summer and fall, galls dry up and open and the nymphs emerge,



Figure 60. Gall of spruce gall adelgid on spruce.

complete the next phase of their life cycle. The galls may persist on the branches for several years.

new shoots, and developing cones. Nymphs are oval and black with a white waxy fringe.

Adults are dark brown and covered with white,

wooly wax. Feeding on

cause galls to form but rather yellowing and twisting of the needles.



Figure 61. Cross section of gall showing nyumphs of spruce gall adelgid on spruce.

Biology: The entire life cycle requires 2 years. There are six stages in addition to the eggs and crawlers when both host trees are present.

Effects: This insect has a very complex life cycle involving two hosts, spruce

galls are unimportant in forest situations. However in nurseries, plantations, and on ornamentals, galls are a concern since they kill branch tips and can stunt and deform trees.

needles turn yellow and become twisted. Heavy infestations on poor sites result in defoliation.

Similar Insects and Diseases: Damage on Douglas-

needle midges or needle cast.

References: 23

Figure 62. Waxy filaments ("cotton") created by Cooley spruce gall adelgid on Douglas-fir branch.

Spruce Aphid Elatobium abietinum (Walker)

Hosts: Engelmann spruce and blue spruce

Symptoms/signs: The spruce aphid is a small green, soft-bodied insect about 1 to 1.5 mm in size. Both winged and wingless forms occur. Spruce aphids feed by inserting their needle-like mouthparts into host foliage and sucking the sap from the needles.

First symptoms of feeding are yellow patches on the needles. If the population increases, discoloration

affected needles turn brown and drop prematurely.

Biology:

Like other aphids, the spruce aphid bears live young with females producing



Figure 63. Adult spruce aphid.

females. Nymphs mature within 3 weeks. During favorable years, large colonies develop during the winter and feed during mild periods. Populations reach a low point during the summer and may

feeding on the current year's foliage. Greatest population increases generally occur from late winter into early spring in northwestern North America and Europe, however, the highest population densities in the Southwestern United States occur in the fall.

Effects:

late spring and early summer, the new foliage is unaffected by the overwintering population of aphids. Current year's foliage is not affected until aphids begin building up again in the fall. Defoliation




Figure 65. Closeup of defoliation caused by spruce aphid.

Pinyon Needle Scale Matsucoccus acalyptus (Herbert)

Host:
Symptoms/Signs: Pinyon needle
cotton looking egg masses in spring, black
bean-shaped nymphs in fall, or emerged

Biology: Adults emerge during middle to late April. Adult females are

adults in April.

winged. After mating, females lay clusters of yellow eggs covered with white, cottony webbing around the root collar, undersides of branches, branch crotches, or crevices in rough bark. Sometimes egg masses are

a rock or log. Crawlers emerge about 5-6 small, dark, bean-shap weeks after eggs are laid. They climb to the ends of branches and settle on the previous year's new growth. After inserting their piercing sucking mouthparts into a needle, they become immobile.




Figure 67. Pinon needles damaged by pinyon needle scale have a chlorotic (yellow) appearance.



Figure 66. Adult pinyon needle scales appear like small, dark, bean-shaped objects on piñon needles.

The body becomes covered with wax and turns black in color. This occurs in the Southwest by early June. By October, the larva resembles a small black bean 1.5 mm long. They overwinter in this stage.

Effects: Infestations can cause needles to yellow and drop. Repeated attacks cause reduced new growth and stunted needles. In severe outbreaks,

small trees may be killed outright and larger trees can be predisposed to attack by bark beetles. Similar Insects and **Diseases:** Symptoms of needle scale may look similar to needle cast caused by fungi, but fruiting bodies will be embedded in needles rather than external. References: 8, 23 Figure 68. Egg masses of pinyon needle scale are found at or near the base of infected piñon. Figure 69. Damaged piñon caused by pinyon needle scale.

Prescott Scale Matsucoccus vexillorum (Morrison)

Host: Ponderosa pine

Symptoms/signs: The most noticeable symptom is the

nodes.


Biology:

and winged males emerge from overwintering pre-adults. Following mating, eggs are laid, often at branch nodes. Eggs are laid in groups,

settle in temporary feeding spots. The second stage larvae settle in permanent feeding locations in July and August. They feed beneath scales at the base of needles and in cracks and crevices of twigs,

or pre-adult stage develops and then overwinters. The life cycle is completed in 1 year.

Affected twigs exhibit lesions, brownish areas in the phloem. A fungus, Cenangium spp., is sometimes associated with the lesions.



Apparently the principal factor in natural control is branch dieback. Few natural enemies have been found outside of a species of ladybird beetle and a green lacewing.

Figure 70. Waxy filaments ("cotton") created by Prescott scale in Arizona.

Effects:

the most common symptom. It occurs primarily in younger trees, less than 75 years old and is most conspicuous in dense stands. Both lateral and terminal shoots can be affected. Mortality rates averaged 1-2 percent in past outbreaks.

Similar Insects and Diseases: Bark beetles known as twig beetles

Figure 71. Branch damage ("flagging") caused by Prescott scale in the White Mountains, Arizona.

also cause twig dieback in ponderosa pine. However upon close

inner bark or the pith of affected branches.

References: 23, 62




Figure 72. Seasonal life history of Prescott scale.

Pine Needle Scale Chionaspis pinifoliae (Fitch)

Hosts:

Symptoms/signs: Damage is especially noticeable on ornamental pine and spruce trees growing along dusty roads. Insects feed by sucking sap from needles, causing the needles to yellow and eventually drop.

Biology: The pine needle scale has two generations per year. The 3.5 mm long mature female scales are most conspicuous. They are almost pure white, slender at the



Figure 73. Closeup of pine needle scale.

front with a wider rear end. Males are smaller and slender and rarely seen. Twenty to 30 eggs are laid in the fall and winter beneath the dead female scale. Eggs hatch in May and the nymphs, or "crawlers", move to new green needles to feed. Nymphs mature by early July, adults mate, and new clusters of eggs are laid. Scales of this second generation mature by fall and lay the overwintering eggs.

Effects: Heavy infestations over several years can kill young trees and severely weaken larger trees, predisposing them to attack by other pests.



Figure 74. Ponderosa pine branch with pine needle scale.

Similar Insects and

Diseases: Damage may be similar to that caused by other agents that discolor needles such as black pineleaf scale, pine needle casts, winter desiccation, drought, and aphids. If scales are present, diagnosis is assured.

References: 8, 32

Ponderosa Pine Resin Midge Cecidomyia piniinopis (Osten Sacken)

Host: Ponderosa pine **Symptoms/signs:**

noticeable in early summer when the new shoots start to droop, the foliage turns yellow and the shoot dies. Larvae are red in color and feed in pits under the bark of the current year's shoots, causing twig deformities. Adult midges



Figure 75. Closeup of ponderosa pine resin midge damage.

resemble mosquitoes. They possess long antennae that are beaded in appearance along with a much reduced wing venation.

Biology: The ponderosa pine resin midge has one generation per year. Larvae overwinter in pits under the bark. In the spring, they make their way to the surface where they spin cocoons on needles.

Effects:

Although yellowing and shoot death

these effects are not considered to be commercially important in most situations.

Figure 76. Larva and pupa of ponderosa pine resin midge.

Similar Insects and Diseases: Twig beetle (*Pityophthorus* spp.) feeding may cause similar death of small twigs and branches on pines.




Figure 77. Flagging damage to branches of ponderosa pine caused by ponderosa pine resin midge.

Pinyon Spindle Gall Midge Pinyinia edulicola (Gagné)

Host:

Symptoms/Signs: Presence of spindle-like swellings at the base of needles that are about 1 cm long with tiny, orange maggots inside.

Biology: Adults lay eggs on needles in late June and early July. Larvae hatch soon afterward and mine into the current year's needles near the base. The plant forms a gall around the feeding larvae. Each gall contains 5 to 40 small, orange legless maggots. The larvae overwinter in the galls and pupate in the spring




Figure 78. Gall on piñon caused by pinyon spindle gall midge.

Effects: This is a common forest insect that rarely causes serious damage. However, heavy infestations in urban settings cause serious defoliation when needles drop prematurely.

References: 8, 23

F C n s,

Figure 79.
Orange-colored
maggots of pinyon
spindle gall
midge.

Spittlebugs Aphrophora spp., Clastoptera spp.

Hosts: Oak, juniper, and southwestern dwarf mistletoe **Symptoms/signs:** The most conspicuous evidence of infestation is the spittle-like froth that surrounds the feeding nymphs.

Biology:

Adults usually lay eggs just under the bark of twigs in summer. Eggs overwinter and hatch in spring. Nymphs feed on twigs and leaves and cover themselves with spittle. The spittle protects nymphs from drying and discourages natural enemies. Nymphs migrate to foliage in July and August where they transform to adults. Nymphs and adults feed on the sap of their hosts via their piercing mouthparts.




Figure 80. "Spit" caused by spittlebug nymphs feeding on pine.

The main natural controls include hot, dry weather that can desiccate

parasites, ants, spiders and birds.

Effects: When these insects are abundant, their feeding can cause discoloration and sometimes mortality of infested twigs.

Similar Insects and Diseases: The juniper twig girdler, *Styloxus bicolor*

may be distinguished from spittlebugs by looking closely at affected twigs. The twig girdler bores inside the twigs, leaving them hollow inside.

References: 23

Spider Mites Oligonychus spp.

Hosts: Conifers and hardwoods **Symptoms/signs:** Spider

mites suck plant juices with needle like mouthparts, which cause spotting, yellowing, fading, and premature drop of foliage. When populations are heavy, cast skins, webbed foliage, eggs, and other mite activity can be found. Adult spider mites are tiny greenish to reddish colored

arachnids with black heads. **Biology:** There can be



Figure 81. Adult spider mites.

the egg stage bear legs.

Effects: Spider mites are mainly an ornamental tree problem and often go unnoticed in forested situations. Outbreaks of spider mites generally last 1 year and are kept under control by natural enemies, such as predatory mites.

Similar Insects and Diseases: Aphid feeding can have the same symptoms of yellowing, spotting or fading foliage. Adult aphids have winged and wingless individuals. In general, aphids are

Figure 82. Damage to juniper caused by spider mites.

larger in size and have three pairs of legs instead of four.

Leaf injury from compounds such as ozone may have similar symptoms such as yellowing, spotting, or fading.

References: 23, 45

ark beetles are one of the most destructive insects in western coniferous forests. It has been estimated that 90 percent of insect-caused tree mortality and more than 60 percent of the total insect-caused loss of wood growth in the United States is due to bark beetles. In the Southwest, bark beetle killed trees were scattered over more than 2 million acres between 2001 and 2003.

The Southwest has a large complex of bark beetles composed of many genera and species. Frequently, several species are found attacking the same host tree and, therefore, it may be


of *Dendroctonus* are the most notorious tree killers in the western United States, *Ips* species also play a very important role in pine forests of the Southwest.

Bark beetles derive their name from their habit of living and mining between the bark and wood of trees and shrubs. Adults excavate egg galleries in bark phloem. All bark beetle life stages are spent in the phloem, inner bark and bark, except when adults leave

beetles feed on the phloem during adult and larval stages.

Most bark beetles are considered secondary mortality agents because they prefer weakened host material. However, during environmental conditions favorable for beetle development, populations may build up rapidly and successfully attack healthy trees. Most bark beetles have a symbiotic relationship with blue-stain fungi. The blue stain fungi can completely penetrate the sapwood within a year. The fungi occlude the outer conducting tissues in the xylem that halts upward water translocation. This action, plus that of the bark beetle feeding, causes the death of a host tree.

Bark beetles produce chemical compounds called pheromones that are used to communicate with other beetles. Aggregation pheromones cause beetles to congregate in certain






Figure 86. Exit holes of bark beetles emerging from

ponderosa pine.

attack trees.
Anti-aggregation
pheromones
cause beetles
to disperse to
neighboring trees
or other areas.
Pheromones
of many bark
beetles have

areas and mass-

and synthetically produced. Both aggregation and anti-aggregation pheromones have been effective to mitigate impacts caused by some bark beetles in the western United States.

Crowns of successfully attacked trees turn from green to yellow to reddish brown. This color




Figure 87. Small rust-colored pitch tubes are a sign that trees have been attacked by bark beetles.

Figure 88. Boring dust in branch crotches, bark crevices or around the base of trees is a sign that trees have been successfully attacked by bark beetles.




Figure 89. Blue stain fungi introduced during bark beetle attacks on conifers occurs in the sapwood.

change, an indication of a dying tree, may occur from a month to more than 2 years after successful attack depending on the temperature, moisture conditions, and density of beetles in the tree. Close inspection of infested tree trunks will show either small globules of resin, small holes through the bark, or reddish boring dust in bark crevices and around the tree base. The removal of bark from infested trees will reveal two types of galleries, egg and

larval. Egg galleries constructed by adult beetles are rather uniform in width. Larval galleries depart at right angles from egg galleries and increase in size as the young grow.

References: 23, 24, 40, 115

Pine Engravers *lps* spp.

Hosts:

Symptoms/signs: In standing trees, fading tops of large trees or whole crowns in small trees can be indicators of Ips infestation. Other external evidence consists of accumulations of boring dust in bark crevices and at the base of the tree. Occasionally pitch tubes can be found on the trunk. Characteristic egg galleries may be found under the bark, slightly engraving the sapwood, hence the common name engraver beetle. In slash, look for boring dust and galleries.

Adults are small cylindrically shaped brown beetles a few millimeters in length. They possess a pronounced concavity at the rear end of the elytra that is bordered on either side by three to six tooth-like spines. Larvae are white, legless grubs.




Figure 90. Egg galleries of Ips pini on fire-damaged ponderosa pine. Note egg galleries radiate outward from centralized nuptial chamber.




Figure 92. Adult Arizona fivespined ips beetle. Note spines along posterior edge of the elytral declivity.

Biology: Beetles produce 2 to 4 generations per year, depending on climate, species, and elevation. In spring, adult beetles emerge from material infested the previous fall and

prefer fresh debris from logging, construction activity or natural events, but during outbreaks living trees may be attacked.

Effects: Bark beetles in this genus have the potential to




Figure 94. Ips species typically initiate attacks near the top of ponderosa pine causing a characteristic top-down fading pattern.



Figure 93. Frass piles indicating Ips attacks on fresh ponderosa pine slash.

kill thousands of pine trees during short-lived outbreaks in Arizona and New Mexico. Pine mortality is frequently noticed in drought years. Pinyon ips (*I. confusus*) killed

across the Southwest from 2002 to 2004. Similarly, the pine engraver beetle (*I. pini*) and the Arizona *I. lecontei*) killed

millions of ponderosa pine during the same time span throughout Arizona. Some outbreaks are associated with human activities that create large amounts of fresh pine debris or that weaken trees. These insects are

providing a food source for other animals.

Similar Insects and

Diseases: Other bark beetles may be found in southwestern pines. They may be distinguished by egg gallery characteristics and adult appearance. Ips egg galleries possess a nuptial chamber—an enlarged excavated area—with one to many galleries radiating from it. The egg galleries are

free of frass. *Dendroctonus* galleries vary by species in shape but lack the nuptial chamber and are packed with frass. Ips adults display a pronounced concavity at the rear end of the elytra, which contains three to six spines on either side. *Dendroctonus* adults on the other hand have a rounded declivity with no spines. Frequently, both *Dendroctonus* and *Ips* species occur within the same tree with *Ips* species typically located in the top half of the bole and *Dendroctonus* in the bottom half.



Figure 95. Stand level ponderosa pine mortality caused by Arizona fivespined ips on Prescott NF, Arizona.





Figure 96. Initial construction of egg galleries by pinyon ips on piñon.

References: 24, 48, 52, 61, 75



Western Pine Beetle Dendroctonus brevicomis (LeConte)

Host: Ponderosa pine

Symptoms/signs: Larval feeding in the inner bark and adult mating and egg laying creates mazelike galleries. Entrance holes, reddish dust in bark crevices, and pitch tubes are indicators of attack. Exit holes are created when the mature beetle leaves the host. Trees, which have been successfully attacked by beetles, are most noticeable when the foliage begins to fade from green to yellow to



Figure 101. Egg gallery of western pine beetle on ponderosa pine. Note that galleries have a crisscrossing, serpentine pattern.




Figure 102. Cross section of bark showing western pine beetle larvae in outer bark.

red to gray. Also, infested trees often exhibit signs of woodpecker feeding activity in the bottom half of the bole. Needles fall off approximately 1 year after the initial attack.

Biology: Adult beetles usually begin

suitable host trees in late spring or early summer and continue until cold weather begins. Females produce from one to three broods a year. Eggs are laid in niches along galleries. When the eggs hatch, the larvae feed in the inner bark working away from the egg gallery.

Effects: The direct effect of successful attack is tree mortality. Usually, the beetles breed

in and kill scattered, overmature, slow growing, decadent, or diseased trees and trees weakened by stand stagnation,

This tree mortality may be considered part of the normal ecological process of succession during which a forest matures and replaces itself. Epidemics can affect ponderosa pine ecosystems by reducing the density and size distribution of its host, and altering species composition and stand structure.

Similar Insects and Diseases: Other bark beetles attacking ponderosa pine. Western pine beetle egg galleries can be distinguished from other beetles Figure 103. Pitch tubes indicating western pine

indicating western pine beetle attack are frequently small in size (<1/4 inch, 6.4 mm).

by *mm*). their maze-like appearance.

References: 17, 23

Figure 104. Woodpecker activity ("flaking off of bark") on ponderosa pine indicating that western pine beetle larvae are in the outer bark.

Mountain Pine Beetle Dendroctonus ponderosae (Hopkins)

Hosts: Ponderosa pine, white pines

Symptoms/signs: External evidence on green infested trees consists of pitch tubes. On successfully attacked trees, these are small and numerous. Pitch tubes on unsuccessfully attacked trees

are larger in size (around 2 cm in diameter), typically white, and widely scattered over the trunk. During drought years, infested trees may not produce pitch, and external evidence consists only of boring dust. These are referred to as blind attacks. Later, foliage on successfully attacked trees fades to yellowish green, to sorrel, red



Figure 105. Adult mountain pine beetle.

Figure 106. Egg larval galleries of mountain pine beetle on limber pine. Note that egg galleries have an initial "crook" or "j" and then run vertically up the bole. Larval galleries run perpendicular to egg galleries.

galleries are straight, vertical, and packed with boring dust.
They range from 10 to 122 cm in length. Adult beetles are brown cylindrically shaped beetles about 4 to 7.5 mm long. Larvae are small white grubs with tan head capsules.

Biology: The beetle produces one generation per year. Adults begin attacking trees in early July and the attack period can extend into September. Females initiate attacks and release pheromones to attract males. Beetles create egg galleries in the inner bark and females lay eggs. In 1 to 2 weeks small larvae

hatch. Generally they overwinter as small larvae. Larvae resume development the following spring and when mature, excavate oval cells in which they transform into pupae and then adults. Adults then bore out and attack new trees. In the Southwest on ponderosa pine, this insect is found primarily on the Kaibab Plateau in northern Arizona and in northern New Mexico. It can be found on white pines throughout the Southwest.

Effects: The direct effect of successful attack is tree mortality. Epidemics can affect ponderosa pine ecosystems by reducing the density and size distribution of its host, and altering species composition and stand structure.

Similar Insects and Diseases: Several other bark beetles may be found in Southwestern ponderosa pine, including other Dendroctonus species and engraver beetles. These may be distinguished from mountain pine beetle by their egg gallery characteristics and adult appearance. Egg galleries of the western pine beetle are maze-like, and those of the roundheaded pine beetle are vertical and slightly sinuous. Galleries of engraver beetles possess a nuptial chamber with one to several galleries radiating out

from it. Engraver beetle egg galleries are free of frass. Ips adults display a pronounced concavity at the rear end of the elytra that possesses three to six spines on either side. The elytral declivity on Dendroctonus adults is rounded and does not possess any spines.


References: 1, 74 Figure 107. Ponderosa pine mortality caused by mountain pine beetle.

Roundheaded Pine Beetle Dendroctonus adjunctus (Blandford)

Host: Ponderosa pine **Symptoms/Signs:**

Reddish brown boring dust is present in bark crevices and numerous pitch tubes are observed on the bark at entrance holes. In heavily attacked decadent trees, pitch tubes are often missing or very small. Later, the foliage fades from green to yellow or bright red, and eventually a dull brown. Woodpecker activity on the trunk is also a good indication that bark beetles have attacked the tree.

Biology: Adult roundheaded bark beetles attack and colonize susceptible trees in October and November. Females initiate the attack producing an aggregation pheromone



drawing large numbers of beetles to the tree. The beetles chew through the bark and excavate a chamber in the moist tissue beneath the bark. Males enter the chamber and mating occurs. Female beetles then excavate horizontal tunnels in the inner bark region for 25 to 50 mm and then longitudinally with the grain an average distance of 300 mm. Eggs are laid individually in niches on alternate sides of the egg galleries, which are packed with frass. The larvae mine across the grain in the inner bark until in the third instar and then bore into the outer bark to complete development.

Effects: Roundheaded pine beetles attack, colonize, and kill trees in the southwestern United States and Mexico. Periodic outbreaks have killed large numbers of ponderosa pine

Figure 109. Pitch tubes caused by roundheaded pine beetle attacks on ponderosa pine.

in southern New Mexico and southeastern Arizona. In 1950, 16,000 pole and sawtimber-sized trees were infested on 2,500 acres near Cloudcroft and 400,000 pole-sized ponderosa pines were killed on 150,000 acres from Mayhill to Ruidoso in 1971. Similar outbreaks occurred during the 1990s in the Sacramento Mountains in southern New Mexico and

southeast Arizona.

Similar Insects:

The roundheaded pine beetle is similar to other *Dendroctonus* beetles, *Ips* beetles, and other members

of the subfamily Scolytidae. *Dendroctonus* beetles are distinguished from each other mainly by their unique egg galleries and, in the

they attack. *Dendroctonus* beetles are distinguished from *Ips* beetles by their rounded posterior ends and galleries, which are packed with frass. In contrast, the posterior ends of *Ips* beetles are slightly concave with and have several to many small spines and the runways (egg galleries) are free of frass.

References: 8, 23, 60

	Figure 110. Ponderosa pine mortality caused by roundheaded pine beetle in the Pinaleño Mountains, Arizona.
Figure 111. Aerial view of ponderosa pine mortality caused by roundheaded pine beetle in the Pinaleño Mountains, Arizona.	

Southern Pine Beetle and Mexican Pine Beetle Dendroctonus frontalis (Zimmerman) and Dendroctonus mexicanus (Hopkins)

Hosts: Chihuahua, Apache, ponderosa pines

Symptoms/signs: Presence of reddish brown boring dust

on the bark at the entry point of the tunnels. In heavily attacked decadent trees, pitch tubes are often missing or so small they can only be seen up close. Later, the foliage begins to fade, turning from green to yellow or bright red, and eventually a dull brown. Woodpecker activity on the trunk is also a good indication that bark beetles have attacked the tree.

Adults are dark reddish brown, cylindrically shaped beetles

approximately
3 mm in length.
Larvae are white,
legless grubs.
The morphology
of southern
pine beetle
and Mexican
pine beetle are
very similar
and, therefore,

Figure 112. Egg galleries of southern pine beetle on Chihuahua pine. Note that egg galleries are "maze-

like" with branching and interconnections.

to tell apart.
Taxonomists use
the seminal rod of

male beetles as the distinguishing characteristic.

Biology: Although little is known of the southern pine beetle biology in the Southwest, this beetle may have 3 to 11 generations in the southeastern U.S. depending on latitude, elevation, and climate. Both southern pine beetle and Mexican pine beetle seem to have only one generation per year in the Southwest.

Adults construct winding galleries in the inner bark, where eggs are deposited in individual niches on each side of the galleries. The larvae mine for a short distance before boring into the outer bark where they pupate. Mexican pine beetle is thought to have a similar biology.

Effects: Southern pine beetle and Mexican pine beetle killed large numbers of Apache and Chihuahua pines throughout 1999-2001 in the Chiricahua Mountains of southeastern Arizona. Pine mortality ranged from single trees to large groups of more than 100. Southern pine beetle has been collected from traps in northern Arizona.

Similar Insects and Diseases: Southern pine beetle and Mexican pine beetle can be distinguished from other pine bark beetles by the shape of egg galleries and beetle morphology.




Figure 113. Chihuahua pine mortality caused by southern pine beetle in Chiricahua Mountains, Arizona.

Red Turpentine Beetle Dendroctonus valens (LeConte)

Hosts: Pines Symptoms/signs:

The best indicators of red turpentine beetle attack on pines are: large pink-white pitch tubes on the lower bole; accumulations of reddish brown sawdust at the base of the tree and in bark crevices; and accumulations of cream to pink colored crystallized resin granules at the tree base. The egg galleries under the bark are fairly wide and linear to




Figure 114. Large pitch tubes at bases of ponderosa pine indicate red turpentine beetle attacks.

irregular in shape. Galleries extend downward from the entry hole 7 cm to 1 m and may even extend into large roots.


Adults are reddish brown, cylindrically shaped beetles approximately 8 mm in length. Larvae are white, legless grubs.

Biology: The number of generations varies from 1 to 2 years

Figure 115. Red turpentine beetle frequently attacks fire-damaged ponderosa pine as indicated by pitch tubes and frass.

in the coldest portions of its range to 2 to 3 generations per year in the warmest areas. Attacks are made in the lower bole of pines. Attacks occur throughout warm weather but peak by midsummer. These beetles often attack trees scarred by

Effects: Through repeated attacks this beetle can sometimes kill trees;



Spruce Beetle Dendroctonus rufipennis (Kirby)

Hosts: Engelmann spruce is principal host, blue spruce is an infrequent host Symptoms/signs:

External evidence on green infested trees consists of entrance holes in the bark and occasional pitch tubes. However, pitch tubes are often indicative of an unsuccessful attack. Red boring dust from entrance holes usually accumulates in bark crevices around the bases of infested trees. Woodpeckers frequently remove large sections of bark looking for larvae to feed on. About 1 year after attack, needles of infested trees usually turn a yellowish green and fall, though some may remain green until the




Figure 118. Egg and larval galleries of spruce beetle. Egg galleries are vertical with eggs laid on alternate sides of the gallery.

Figure
119. Fading
Engelmann
spruce mortality
caused by
spruce beetle
in White
Mountains,
Arizona.

second year. Needles do not turn a reddish color like those on bark beetle infested pines. Under the bark, egg galleries are vertical and slightly groove the xylem. Egg galleries range from about 6 to 30 cm




Figure 120. Engelmann spruce mortality caused by spruce beetle on the Pinaleño Mountains, Arizona.

long. Adult beetles are cylindrically shaped and 4 to 6 mm long. They are generally dark brown to black with reddish brown wing covers; however, older adults are usually entirely black. Larvae are creamy white, cylindrical, legless grubs. Larvae reach a length of about 6 mm. Pupae are creamy white and similar in size to the adult.

Biology: Spruce beetles generally produce

one generation in 2 years; however, generation time can take from 1 to 3 years. Adults emerge from May through July and begin attacking new host trees. Eggs hatch and larvae develop during the summer. Broods overwinter as larvae or callow adults and complete development to mature adults by the following August. Spruce beetle attacks standing trees, windthrown trees, or logging residuals. It prefers large diameter, green downed material to standing trees. Populations are affected by a number of factors including the number of susceptible hosts, natural enemies, and extreme cold.

Effects: The direct effect of spruce beetle attack is tree mortality or in some cases strip kill, when only a vertical section of the bole is attacked. Strip attacked trees often die in subsequent years. Spruce beetle epidemics can have both immediate and long-

changes in stand structure and composition. The longer-term effects come from successional changes that result from microclimate changes as well.

Similar Insects and Diseases: The most common associates are in the genus *Ips*. In New Mexico, *I. pilifrons sulcifrons* is the most common, while in Arizona, *I. pilifrons utahensis* is found most often. *Ips* beetles frequently infest the upper (exposed) portion of downed material, while spruce beetle favors the shaded material. *Ips* egg galleries differ from those of spruce beetle by containing a nuptial chamber and by being free of frass.

References: 41, 86



Figure 121. Flaking of bark caused by woodpeckers indicates spruce beetle larvae in the outer bark of Engelmann spruce.

Douglas-fir Beetle Dendroctonus pseudotsugae (Hopkins)

Host:

Symptoms/signs: Reddish-

been attacked. At times, the most evident sign of attack is the clear resin exuding from entrance holes on the upper portions

always present on successfully attacked trees. Needles of successfully attacked trees change from green to yellow to sorrel to reddish brown in the year following attack. Egg galleries run parallel to the wood grain and are 20 to 25 cm long. Adult beetles are stout, cylindrically-shaped, and 4 to 6 mm long. The head

and thorax are black; the wing covers reddish brown,

becoming darker with age.

Biology:

has one generation per year. Although

Figure 122. Egg and larval galleries of

Douglas-fir beetle. Note

larval galleries occur in

alternating groups.

new attacks occur in late spring to early summer. Broods develop under the bark throughout the summer and early fall. Adult, pupal, or larval life stages can overwinter. Brood that develops to pupae or adults by winter usually emerges between April and June. Overwintering larvae emerge as adults in summer. A small percentage of adults that overwinter will re-emerge from the spring-attacked

Figure 123. Douglas-fir beetle caused mortality of dwarf mistletoe-infected Douglas-fir on the San Francisco Peaks. Arizona.




Figure 124. Frass (boring dust) in bark crevices, spider webs, or at base of Douglas-fir indicates successful attack by Douglas-fir beetle.

trees and attack additional trees in the middle of summer. Typically, late-season attacks account for less than 20 percent of all attacks in one season.

Effects: The direct effect of successful attack is tree mortality. At low or endemic levels, the beetle infests scattered trees that are infected with dwarf mistletoe and/or armillaria root disease, injured

agents. During outbreaks, groups of 100 or more attacked trees are

long-term effects on mixed conifer forests. Immediate effects include changes in stand structure and composition. Successional (longterm) changes result from alterations in microclimate.

Similar Insects and

Diseases: Other bark beetles,
Scolytus monticolae and Douglas(Pseudohylesinus
nebulosus), often occur in large
branches, boles of pole-sized
trees and the top of trees killed by

separated by gallery patterns and characteristics of adult beetles.

References: 23, 89




Figure 125. Galleries of Douglas-fir pole beetle on small diameter Douglas-fir. Galleries are short, longitudinal and often with two branches.

Western Balsam Bark Beetle Dryocoetes confusus (Swaine)

-		
ш	Oct :	
ш	wst.	

Symptoms/signs: Small entrance holes and boring dust

may be evident. Under the bark, egg galleries lightly etch the surface of the wood. Egg galleries radiate from a central nuptial chamber. Attacked trees generally turn yellow-red within a year. Needles on a

dark brown, cylindrical beetles about 2 to 5 mm long. The posterior end of the elytra is abruptly rounded with no spines.

Biology:
Little is published on the biology of this species. A 1- to 2-year life cycle

on the biology of this species. A 1- to 2-year life cycle is indicated. The lower to mid bole of standing trees is usually selected for attack. This

Figure 126. Egg gallery of western balsam bark beetle on corkbark fir. Egg galleries typically have five or more galleries radiating from a nuptial chamber.

Figure 127. Adult western balsam bark beetle. Note the "crew cut" appearance on front of beetle heads.


species is moderately aggressive and capable of killing older and/or weakened trees. Occasionally cut or fallen trees are attacked.

Effects: The direct effect of successful attack is tree mortality.

Similar Insects and Diseases: Other bark beetles

species of *Scolytus* and *Pseudohylesinus*. They may be separated by gallery patterns and characteristics of adult beetles. Western balsam bark beetle attacks are frequently associated

References: 23, 64, 115



Fir Engraver Scolytus ventralis (LeConte)

Hosts:

Symptoms/signs: Freshly attacked green trees have tiny holes in the bark, reddish-brown boring dust in bark crevices, and

streams of clear pitch exuding from entrance holes. Needles change from green to yellowgreen to sorrel and then red to brown. Larvae, pupae, and egg are found in the inner bark; egg and larval galleries deeply score the wood, creating characteristic horizontal egg galleries and vertical larval galleries. Adult beetles are small, shiny, black beetles about 4 mm long. The abdomen is incurved at the rear.

Biology: In warm locations, the




Figure 129. Egg and larval galleries of fir engraver. Galleries are horizontal with two branches radiating out from entrance point.

one generation and a pa	artial second generation each year. In cooler
	sites, the beetle needs 2 years to
	Figure 130. Adult fir engraver
	Figure 130. Adult fir engraver beetle. Note the "sawed-off"
	appearance of the beetles,"
	abdomen.

occurs throughout the summer, but peaks in July and August. Fir engravers overwinter primarily as young or mature larvae. In the spring, they feed for a short period and then construct pupal cells at the end of their galleries.

Effects: Fir engravers are a major mortality agent of true

pole-sized to sawtimbersized trees. Outbreaks often occur during and following droughts. Root diseases are often associated with attacked

can also breed in slash and wind-thrown trees. Figure 131. White fir mortality caused by fir

engraver on Mormon Mountain, Arizona.

only partially attacked.

Similar Insects and Diseases: Other bark beetles attack Dryocoetes and Pseudohylesinus.

They may be separated by gallery patterns and characteristics of adult beetles.

References: 3, 19, 23

Twig Beetles Pityophthorus spp., Pityogenes spp., Pityoborus secundus

Hosts: Ponderosa	
Symptoms/Signs:	
Trees attacked and colonized	
by twig beetles exhibit fading	
needles on twigs and branches,	
and twig and branch dieback	
hroughout the crown. Tan	
awdust is produced around	
he attack sites. On smaller	
wigs and branches, most of the	
cambium is mined beneath the	
oark. Small, star-shaped egg	
galleries generally occur under	
he bark of the larger branches	
and small trunks.	
Piology Adult twice	

Biology: Adult twig beetles are about 3 mm long and dark brown. Most species

Figure 132. Pitch tubes of twig beetles on piñon are typically very small (<1/8 inch).

Figure 133. Damage to ends of piñon branches caused by twig beetles. Very small pitch tubes may be present on twigs.

have a rounded rear end, but a few have a pair of short spines. The larvae are fat, white, C-shaped grubs with light brown head capsules and feed under the bark. Most species have 2 to 4 generations per year, depending on local conditions.

Effects: Typically attacks and kills small twigs and branches of drought stressed or otherwise weakened pines and other

conifers. Although twig beetles are generally considered of secondary importance, under favorable conditions they may

high numbers and attack and kill small trees. *Pityogenes carinulatus* killed thousands of young ponderosa pine in 1967 in New Mexico as a result of populations increasing in thinning



Figure 134. Mortality of piñon regeneration killed by twig beetles near Los Alamos, NM.

and logging slash. Twig beetles infested high numbers of young

Similar Insects and Diseases: See *Ips* discussion. Most twig beetles have elytral declivity and egg galleries that are somewhat similar to *Ips*; however, twig beetles are smaller in size and are typically found in branches and very small diameter trees.

References: 9, 23

Cedar Bark Beetles Phloeosinus spp.

Hosts: Arizona cypress and junipers

Symptoms/signs: External evidence consists of twig killing

are simple and longitudinal, 2 to 7 cm long, usually engraving the wood rather deeply. Egg niches are usually rather large and conspicuous. Larval galleries wander away from the parent galleries.

Adults are reddish brown to black, shiny beetles ranging in size from 2 to 4 mm long. Larvae are small white grubs, with brown head capsules.

Biology: One to one and a half generations are produced per year. Attacks occur in spring and summer. Adults and larvae feed in the inner bark in galleries. Newly emerged adults feed on the pith of twigs of living trees prior to constructing egg galleries. Often twigs are hollowed out completely and killed.




Figure 135. Egg and larval galleries of cedar bark beetle. Egg galleries have a "thermometer" or "centipede" shaped pattern.




Figure 136. Juniper mortality caused by cedar bark beetle near Flagstaff, Arizona.



Figure 137. Pitch tubes on Arizona cypress caused by cypress bark beetle near Sedona, Arizona. Note that pitch tubes are frequently absent.




Figure 138. Exit holes of cypress bark beetle on Arizona cypress near Sedona, Arizona.




Figure 139. Arizona cypress killed by cypress bark beetle near Sedona, Arizona.

Effects: Main effects on their hosts are twig killing or tree mortality; however, these insects are not aggressive and are generally found attacking trunks, tops and limbs of weakened, dying or felled trees. Occasionally outbreaks occur.


Similar Insects and Diseases:

This is the main bark beetle genus attacking cypress and juniper. There are some woodboring beetles that attack these trees. Bark beetles are distinguished from woodborers by the shape and location of the galleries and size of the adults and larvae. Galleries of woodborers extend both in the bark and wood. Woodborer larvae and adults are larger than bark beetles.

References: 23, 81

Figure 140. Extensive mortality caused by cypress bark beetle north of Clifton, Arizona.

his group contains members of several orders (Lepidoptera, Hymenoptera, and Coleoptera) that bore into the sapwood and sometimes the heartwood



Wood Borers Roundheaded borers, Cerambydidae Flatheaded borers, Buprestidae

Hosts: Conifers and hardwoods

Symptoms/signs: Larvae are found under the bark of dead and dying trees. These galleries are usually wider than bark beetle galleries and vary in diameter as the larva grows, in contrast to the egg galleries of bark beetles which have a uniform diameter throughout. Roundheaded borer galleries are packed with a coarse

boring dust. In many species, larvae complete their development in the wood so tunnels extend into sapwood and sometimes the heartwood. Adults emerge through large emergence holes in the bark.

Adult roundheaded borers are medium to large sized, oblong to cylindrical insects. The antennae are often longer than the body, giving them the name "longhorned beetles." Larvae are relatively large, particularly when fully developed, white, cylindrically shaped grubs. The heads are slightly larger in diameter than the body.

compact, and often brightly colored beetles. Larvae have a very

body segment behind the head is much broader than the following body segments and has horny plates on the top and bottom.

Effects: These insects primarily attack weakened, dead or dying trees. A few species attack and kill apparently healthy trees, such as the western cedar borer on junipers and *Agrilus* species on *Populus*. They often attack





Figure 143. Adult longhorned beetles (adults of roundheaded wood borers). Note that antennae are longer than the length of the body.

trees already infested with bark beetles and sometimes compete with them. Larval mines penetrate both the cambial region as well as the wood, sometimes mining it extensively. They are important in the process of nutrient cycling by assisting in breaking down woody



wood borer). Note shorter than the body.

Juniper Borers Trachykele blondeli (Marsuel) Callidium spp Atimia spp Chrysobothris spp Styloxus bicolor (Champlain and Knull)

Hosts: Junipers and Arizona cypress

Symptoms/signs: Exit holes.

Effects:

are aggressive pests in drought-stressed junipers and cypress. Damage can be extensive before symptoms are apparent. Usually a large portion of the tree or the entire tree dies before the insects' exit holes are noticed.

Biology: *Callidium* spp. larvae bore beneath the bark making very wide tracks that distinctively score the outer sapwood much like a router. Older larvae excavate oval tunnels deep in the wood and overwinter. Adults emerge throughout the warm months of the year. There is one generation per year. Adult beetles are rather short-horned for cerambycids and dark blue or black. These roundheaded borers leave an oval or rectangular exit hole.

Atimia spp. are small roundheaded borers about 6.5 mm




Figure 146. Larval feeding galleries of the western cedar borer.

long and generally have a 1-year life cycle. These longhorned beetles attack thin bark portions of seriously weakened and dead juniper and cypress.

Trachykele blondeli, the western cedar borer, is 11 to 17 mm long and attacks juniper and Arizona cypress. Females lay eggs under bark scales and branches of living

trees. Flatheaded larvae bore from the branches into the main bole. They feed primarily in the heartwood for several years. Adults emerge in the spring, leaving oval or rectangular exit holes.

for several years. Adults
emerge in the spring, leaving
oval or rectangular exit
holes.

Chrysobothris

Figure 147. Larval feeding galleries and
exit holes of the western cedar borer.

cypress. They are mediumsized beetles about 11.5 mm

long. Larvae bore into the bark and outer wood of weakened trees.

The juniper twig pruner, *Styloxus bicolor*, is a small roundheaded borer that mines the small branches and causes twig dieback during dry periods.

Similar Insects and Diseases: See cedar bark beetles.

References: 9, 23

Figure 148. Juniper damage caused by western cedar borer.

Ambrosia Beetles Trypodendron spp, Platypus spp, Gnathotrichus spp, Xyleborus spp

Hosts:

maple

Symptoms/signs: Entrance holes are small for most species (1.5 mm or less) and are marked by piles of white boring dust. Gallery patterns vary by species, but all tunnels run perpendicular to wood grain. The mines of *Platypus* species penetrate into the heartwood in contrast with most ambrosia beetles in the Scolytidae that mine only the sapwood.

Effects:

Weakened, dying, recently cut or dead trees are attacked. Galleries cause defect in logs. Some species can attack freshly cut lumber before it has been dried. Populations can build up in

killed trees, bases of trees attacked by bark beetles, logging slash, and logs in storage. These beetles are Figure 149. White boring dust caused by ambrosia beetles tunneling in the wood of a bark beetle-killed pine.

unique among bark beetles in that larvae feed upon a special type of fungus, known as the ambrosia fungus, which grows in the galleries. Larvae do not feed on wood or phloem as in other bark beetles.

Biology: Insect appearance varies between beetles. Adult *Trypodendron*, *Gnathotrichus*, and *Xyleborus* beetles are all dark colored, cylindrically shaped beetles, about 2 to 4.5 mm

long (depending on the species). Larvae are white, legless grubs with brown head capsules. All three genera belong to the family Scolytidae. *Platypus* on the other hand belongs to the family Platypodidae, which is closely related to the Scolytidae. Adult *Platypus* are more elongate, cylindrically shaped beetles than scolytids.

Similar Insects and Diseases: May be confused with other bark beetles; however, ambrosia beetles are the only ones that bore straight into the bole

dust.

References: 23, 32

Figure 150. Entrance points of adult ambrosia beetle. Holes are stained by blue stain fungi.

one and seed insects, also called conophytes,

Destruction of forest seeds may be caused by

varies dramatically from year to year and by location, so do impacts caused by cone and seed insects. In general, cone and seed insects are relatively unimportant under forest conditions as the amount of seed produced by forest trees usually far exceeds the number of seeds destroyed by these insects. However, they can have economic impacts to seed orchards in which seed production is limited. They can also impact natural regeneration of white pines that are being severely impacted by white pine blister rust. The most injurious groups of cone and seed insects in the Southwest are the seed moths and coneworms (Lepidoptera), seed worms and midges (Diptera), and cone beetles (Coleoptera).

References: 5, 23, 36, 47, 87

Cone Beetles Conophthorus spp.

Hosts: Pines Symptoms/signs:

External evidence consists of pitch tubes at the point of entry on the cone stalk or the base of the cone. Following attack, cones turn brown and wither. If beetles have emerged, small round exit holes may be found on the outside of affected cones. Inside, beetles leave cones riddled with tunnels and frass. Adult beetles are reddish brown to black, shiny bark beetles. Larvae are small white, legless grubs.




Figure 151. Adult boring hole indicating attack on ponderosa pine cone by Conophthorus cone beetles.

Biology: One generation is produced per year. Adult beetles bore into the base or stem of immature second-year cones in the spring. A gallery is created along the cone axis, with eggs deposited along its sides. The creation of the gallery severs the conductive tissues of the cone, killing it. Larvae hatch and feed on the scales, seeds, and tissues of the cone. The brood complete development




Figure 152. Damage to interior ponderosa pine cone caused by larva (arrow) of cone beetles.

during the summer within the cone, and usually overwinter there. Some new adults emerge and may bore into shoots or conelets and overwinter.

Effects: These beetles cause the death of second-year cones. The amount of mortality is highly variable but in some years a large proportion of cones may be attacked.

Similar Insects and Diseases: This insect is best distinguished from other insects that attack developing cones by the presence of the pitch tube on the cone base or stem.




Figure 153. Life cycle of cone beetles.

References: 23, 36

Ponderosa Pine Seedworm Cydia piperana (Kearfott)

Host: Ponderosa pine
Symptoms/signs: There is
no external evidence of infestation.
Inside the cones, frass and mining
can be seen in the cone axis. Seed
pairs on the same scales can become
fused together by the silk lined
tunnels produced by the larvae.
Often these seeds tend to stick to the
scales and remain in the cones.

Adult moths have a wingspan of 10 to 20 mm. The fore wings are metallic gray with silver bands. Larvae are 10 to 15 mm when mature, white to cream colored, with a mottled head capsule.

Biology: One generation is produced annually. Eggs are laid at the base of cone scales, on the surface of the scale, or on the cone stalk. Newly hatched larvae bore




Figure 154. Larvae of ponderosa pine seedworm.




Figure 155. Damage of ponderosa pine seed by ponderosa pine seedworm

with frass. It then moves on to another seed leaving a silk lined trail. As the larva matures, it burrows into the cone axis and overwinters. In spring, the insect pupates. Following pupation, the adult emerges.

Effects: In some years this insect can consume a large proportion of the seed crop.

Similar Insects and Diseases: A number of insects affect cones. This insect is distinguished by the presence of the frass packed seeds, larvae or mining in the cone axis.

References: 36

Pine Seed Chalcid Megastigmus albifrons (Walker)

Host: Ponderosa pine

Symptoms/signs: Adults are antlike in appearance and range in color from yellow to black or brown. Females have a long, curved ovipositor. The larvae are small, white and legless. The outer seedcoat develops normally and shows no evidence of an oviposited egg. When adults emerge they leave a small round hole in the seedcoat.




Figure 156. Larva and seed damage caused by the pine seed chalcid.

Biology: Adult emergence

lasts for about a month in spring. Females may mate or reproduce parthenogenetically. Eggs are oviposited through the young scales of developing cones directly into the seed. Usually one egg is laid per seed. The larva matures in the seed, consuming its entire contents. The insect overwinters in the seed and pupates in early spring. Some adults emerge at this time while others may diapause for up to 3 years.

Effects: Pine seed chalcid can cause a large percentage of cull seed in commercial seed production operations. Since it is very

unknowingly transported.

Similar Insects and Diseases: There are a variety of cone and seed feeding insects in the West. Larvae of the family Cecidomyiidae (Diptera) may cause similar reductions in seed production; however, the larvae are not voracious feeders and damage to cones and conelets results from gall formation on the seed scale which prevents seed dispersal. Other species of insects that feed in seeds and cones of ponderosa pine in the Southwest include ponderosa pine cone beetle (*Conophthorus ponderosae*), pine coneworm (*Dioryctria auranticella*), and the ponderosa pine seed moth (*Laspeyresia piperana*).

References: 3, 14

Pine Coneworm Dioryctria auranticella (Groté)

Host: Ponderosa pine

Symptoms/signs: Entire cones are usually killed by pine coneworms; partially killed cones become distorted and do not open.

Biology: Larvae begin feeding in ponderosa pine cones in late spring. They make an entry hole in the basal portion of the cone and consume seeds and scale tissues. Mature larvae pupate in the cavities in the cones created by feeding activity of the larvae. Adults emerge in middle to late summer.

Effects: This obligate seed and cone insect can be destructive when populations are high. Up to 80 percent of cones can be damaged per year in the Southwest. Coneworms feed on seeds and scales from a cavity inside the cone, severely distorting the cone and preventing extraction.

Similar Insects and Diseases: There are a variety of cone and seed feeding insects in the West. Other species of insects that feed in seeds and cones of ponderosa pine in the Southwest include pine seed chalcid (*Megastigmus albifrons*), ponderosa pine cone beetle

(Conophthorus ponderosae), and the ponderosae), and the ponderosa pine seed moth (Laspeyresia piperana). This conophyte is distinguished by the entry hole in the basal portion of the cone and larval feeding cavities



webbing. Figure 157. Damage to ponderosa pine cone caused by larva of the pine coneworm.

References: 5, 87

Bud and Shoot Insects

njury to buds and shoots may be caused by caterpillars, weevils, bark beetles, midges, aphids, or scale insects. Some of these insects also feed on other plant parts such as foliage, cones, and branches. By far the most frequently observed damage to pine shoots is caused by tip moths and shoot borers

Shoot insects affect trees by damaging terminal or lateral shoots. Such damage typically reduces height growth and deforms the tree. Forked branches or multiple leaders often result from a previous year's insect damage. Seedlings and saplings are most susceptible to damage by shoot insects. Trees above 2 m in height

from growing taller than herbaceous competitors, resulting in mortality.

Most of these shoot insects are important in young, intensively managed stands or replanted forested areas. The susceptible stage of hosts is predictable and managers can plan accordingly to minimize impacts caused by shoot insects.

References: 23, 40, 43, 108

Pine Tip Moths Rhyacionia spp.

Host: Ponderosa pine

Symptoms/signs: Insect appearance varies between species.

R. bushnelli adults have a wingspread of 10 to 15 mm; forewings

are mottled yellowish gray and reddish brown. Larvae are yellowish with black heads, and about 9 to 12 mm long when mature. *R. neomexicana* adults have a wingspan of about 0.24 mm. Forewings are irregularly banded with transverse bars of dark gray, blackish and brick-red scales on the inner two-thirds of the wing. Larvae




Figure 158. Adult pine tip moth.

Figure 159. Damage of ponderosa pine shoot caused by pine tip moths.

are usually orange in color with dark brown to light tan head capsules and are about 12 to 16 mm long when fully developed.

Biology: Larvae mine inside new shoots in spring and early summer, killing them. Both lateral and terminal shoots are attacked. Larvae initially mine needles and then bore into the shoot. Pitch tents, frass, and silk webbing are all signs of tip

moth activity. Once in the shoots, larvae feed between the pith and the bark, eventually hollowing out the shoot.

Effects: Larvae mine in the phloem and xylem of lateral and sometimes terminal shoots. Small trees, less than 2 m are most susceptible. Shoots are killed by the attacks. Repeated attacks slow growth and cause crooks, forks, multiple stems, and spike tops. Tip moths rarely kill established trees outright, but attacks can affect survival of young planted seedlings.

References: 23, 43

Similar Insects and Diseases: See western pineshoot borer.




Figure 160. Closeup of damaged ponderosa pine shoot caused by pine tip moths.

Western Pineshoot Borer Eucosma sonomana (Kearfott)

Host: Ponderosa pine

Symptoms/signs: Adults are moths with coppery-red forewings marked with two bright gray transverse bands and a

wingspread of about 20 mm. External symptoms include shortened needles and stunted terminal growth. Following emergence of the larvae, exit holes can be found near the center of the shoot. In the year following infestation, a distinct swelling can be seen around the stem. near the exit hole. Occasionally terminals break off where the stem is weakened by the exit hole.



Figure 161. Damaged (top) and undamaged (bottom) shoots of ponderosa pine caused by western pine shoot borer larvae. Note "bottlebrush" appearance of damaged shoot.

Biology: Eggs are laid on elongating shoots in spring. The larvae feed in the pith of these shoots, vacate in late spring, and drop to the ground to pupate. There they remain dormant through the winter and hatch as adults the following spring.

Effects: Larvae mine only in the pith of the shoot. Usually shoots remain green for at least a year after attack, however, in some cases shoots are killed outright. Most often shoots live, but become stunted. Frequently, lateral shoots overtop affected shoots and a crook or a fork results. The main result of infestation is reduced

height growth, about 25 percent of one year's vertical growth per attack. Depending on management objectives, this may or may not be important. The insect occurs at low population levels.

Similar Insects and Diseases: See pine tip moths.

References: 23, 94, 97

Figure 162. Larva of the western pine shoot borer. Note that larval feeding is restricted to the pith region of the shoots.

Pinyon Pitch Nodule Moth Petrova arizonensis (Heinrich)

Host:

Symptoms/Signs: Attacks are characterized by fading branch tips and small nodules of pitch formed at the insects feeding sites. The pitch nodules are hollow balls of pitch 13 to 25 mm long, round, smooth, and light purple to red in color. Pitch nodules usually occur at the crotch of two or more twigs.

Biology: Small, rusty-brown moths emerge through holes in the pitch nodule in late June and early July. Eggs




Figure 163. Damage caused by the pitch nodule moth.

are laid on the needle sheaths of the current year's foliage. Newly hatched larvae feed on the young needles before boring into the bark at nodes or whorls of twigs or branches. Fullgrown larvae are about 13 mm long, reddish-yellow in color with a black head and a dark area behind the head. Pupation occurs inside the pitch nodule in June. The pupae move just below the pitch surface before they emerge as adults.

Effects: Attacked twigs and branches are killed. Leaders are occasionally damaged and forked trees may result.

Figure 164. Cocoon of pitch nodule moth in damaged ponderosa pine shoot.

References: 9, 23

Bark Moths Dioryctria spp.

and

Pitch Moth Synanthedon spp.

Hosts:

Symptoms/signs: Pitch moth attacks appear as large, ugly

masses of pitch that form at the wound site. Bark moth attacks typically produce less pitch.

Effects: Larger branches, limbs, and trunks of young trees are attacked. Repeated attacks can seriously weaken and kill branches. The most severe damage is to trees under 6 m, especially in urban areas. The insects are rarely a problem on larger trees or in the forest environment.



Figure 165. Damage to piñon caused by bark moths (Dioryctria) at Sunset Crater NP, Arizona.

Biology: Pitch moths require 2 years for one generation to mature, overwintering as larvae each winter. Bark moths require only 1 year for a generation to mature, overwintering as eggs or larvae. Eggs are laid in bark crevices or near mechanical wounds on the bark. Newly hatched larvae tunnel under the bark forming

irregular g sapwood. tree produ Oozing pi cover entr destructiv 15 to 25 n green, or 1 on the inn marked w

Figure 166. Damage caused by pitch moth.

irregular galleries or elongated gouges in the sapwood. Pitch moth larvae feed on pitch the tree produces in response to their tunneling. Oozing pitch masses 25 to 75 mm in diameter cover entry holes and conceal larvae and their destructive tunneling. Full-grown larvae are 15 to 25 mm long, dirty white, yellow, orange, green, or light brown. Bark moth larvae feed on the inner bark and when full grown, are marked with rows of dark spots.

References: 9

Juniper Twig Pruner Styloxus bicolor (Champlain and Knull)

Hosts: Juniper and Arizona cypress Symptoms/ **signs:** The juniper twig pruner adult is 7 to 11 mm long and has a reddishorange head and brownish to black body. Larvae are small, white, and Figure 167. Damage to junipers caused by the cylindrical, legless juniper twig pruner. grubs. **Biology:** The life cycle may take as long as 2 years to complete. Eggs are laid on branches, often near an intersection of twigs, 0.5 m from the branch tip. Grubs kill twigs by boring through the centers. **Effects:** Juniper twig pruner causes twig dieback on Figure 168. Closeup of damage to juniper shoots caused by the junipers and cypress growing juniper twig pruner. throughout the Southwest. Similar Insects and **Diseases:** Cedar bark beetles and western cedar borer. **References:** 9, 42 Figure 169. Exit hole of

juniper twig pruner.

variety of insects attack trees that are recently dead or dying. Losses from these insects are about 1 to 5 percent of the annual timber harvested. Some of these insects have previously been discussed (e.g., woodborers and ambrosia beetles) and others are described on the following pages.

Insects of wood products can either attack moist or dry seasoned wood. Those that attack moist, seasoned wood are usually symbiotically associated with protozoa, fungi, and bacteria that aid the insects in the digestion of cellulose and lignin. Some of these insects will continue to develop in the wood for several years after the wood has been dried and manufactured.

Some of these insects do not obtain their food from the wood. They tunnel into the wood to secure a sheltered base for foraging expeditions and a nursery for the young. Those insects attacking dry, seasoned wood also live in a symbiotic relationship with other microorganisms. These insects can breed successfully in wood with water content as low as 6 percent.

References: 14, 23

Carpenter Ants Camponotus spp.


Hosts:

Symptoms/signs: Carpenter ants are large in size, about 5-15 mm long, and black or black and red in color. They tunnel in the wood of stumps, logs, dead standing trees, the interior dead sections of livings trees, and the wooden portions of buildings. These galleries are honeycomb-like in appearance; the walls are sandpaper smooth, free of frass, and run across the grain. The wood borings,

accumulate at the base of inhabited trees.

Biology: Mating

in the spring. The female then lays eggs, which take approximately 2 months to become mature larvae. The larvae then pupate in cells. Each individual will emerge as one of three possible castes: winged males, winged females, or workers. Carpenter ants do not eat wood but excavate galleries for nesting purposes. These insects are general feeders, and will eat animal food (such as caterpillars, meat, or refuse) and sweets (from aphid honeydew, ripe fruit, or sap.)



pine caused by carpenter ants.

Effects: The excavations in wood can be so extensive that the structural integrity of the tree is lost, predisposing the tree to wind breakage.

Similar Insects and Diseases: Horntails or "woodwasps" (Siricidae) have similar excavations in host trees. The main differences being that they actually have the wood borings pass through the digestive tract and are packed in the tunnel behind the larva. They usually attack fresh logs or newly dead trees. Adults are generally thick-waist, wasp-like insects that are metallic blue or black in color. Adults emerge in late spring or summer and lay eggs in solid wood (conifers or hardwoods) and the life cycle can range from 1 to 2 years. Pupal cells are found in the phloem and the exit holes from mature adults are round.




Figure 171. Cross section of stump with carpenter ant galleries.

Termites Order Isoptera

Hosts: Most tree species

Symptoms/signs: Termite damage is honeycomb in shape and follows the grain of wood. There are three types:

Drywood termites. Coarse sand-like fecal pellets found

Dampwood termites. Typically colonize dead and down trees and untreated wood in contact with the ground.

Subterranean termites. Presence of mud shelter tubes on wood, walls and tree trunks.

Biology: Termites are eusocial insects that have well developed caste systems. Caste consists of primary reproductives,

supplementary reproductives, workers, and soldiers. Only the reproductives have wings and they

swarms outside the colony. As their name implies, workers perform most of the work for the colony. They are generally pale in color, lack



Figure 172. Castes (queen, soldier and worker) of termites.

compound eyes, and have small mandibles. Soldiers are sterile adults with greatly enlarged and armored heads and mandibles. Their sole duty is defense of the colony against invaders. The conversion of cellulose to smaller units by protozoa and bacterial organisms in the guts of termites allow them to digest cellulose from wood and other plants.

Effects: Termites can be extremely destructive to structures built out of wood. Even houses with a cement block foundation are at risk because subterranean termite tunnels can cross over the cement to wood framing. Damp wood termites attack wood that has high moisture content and do not require contact with the ground.

Similar Insects and Diseases: Termites are sometimes referred to as "white ants"; however they can be differentiated from ants by their soft body, light color, straight beadlike antennae, and broad connection between the thorax and abdomen. Ants are hard bodied, darker in color, elbowed antennae, and have a narrow "waist."




Figure 173. Damage to sign post caused by termites in Oak Creek Canyon, Arizona.

Hardwood Leaf Diseases

Leaf diseases in Arizona and New Mexico are most often observed on aspen, Arizona sycamore, cottonwood and willow. Although clonal susceptibility to foliage pathogens is common for aspen and willow; whole areas can become infected during favorable moisture periods. Severe infection can cause premature defoliation and reduced tree growth.




Figure 174. Foliar diseases of aspen occur with adequate spring moisture.

Black Leaf Spot Marssonina populi (Lib.) Magnus

Hosts: Aspen and cottonwood

Symptoms/signs: Small brownish spots appear on infected leaves in late July and early August. The spots later enlarge and turn blackish, are of various sizes and irregular in outline, and have a yellowish to golden border. Infected leaves are often smaller than normal and fall prematurely.

Biology: Primary infection occurs soon after leaves emerge in spring, from spores produced on twig lesions or from infected fallen leaves. Additional spores are produced on newly infected leaves, which initiates the repeating cycle of disease that continues with wet weather until leaves fall. Symptoms intensify and seem to ascend trees as the season advances. *Marssonina populi* survives the winter as tiny stromata in fallen leaves and twig lesions.

Effects: Black leaf spot is the most common leaf disease of aspen in the West. Severe outbreaks may cause foliar browning in midsummer and nearly complete defoliation by early August. Regrowth follows in late summer and early autumn, and twig dieback may follow in winter because late season shoots lack normal cold hardiness. Defoliated trees may folTis onwoay Trns omspoal

Ink Spot Leaf Blight Ciborinia whetzelii (Seaver) Seaver

Hosts: Aspen and cottonwood

Symptoms/signs: *Ciborinia whetzelii* forms brown to black spots, or stromata, in blighted leaves. Infected leaves turn brown and the ink spots, or sclerotia, begin to drop out leaving a circular hole in the dead leaves.

Biology: *Ciborinia whetzelii* infects young leaves in the spring by ascospores produced on fruiting structures that overwintered on fallen leaves. Infected leaves begin to die midsummer, but defoliation may not take place until autumn.

Effects: Ink spot leaf blight is more severe on smaller aspen

trees and in the lower crowns of larger trees. As with many aspen diseases, some clones appear to be more susceptible than others.

Similar Insects and Diseases: This fungus may be confused with *Marssonina populi*; however, *C. whetzelii*

and slightly raised fruiting structures, while those of *M*. *populi* are irregularly shaped

Reference: 38

Figure 176. Black Ciborinia whetzelii fruiting bodies.

Melampsora Rust *Melampsora* spp.

Hosts:

Symptoms/signs: Powdery yellow to yellow-orange spores produced on both sides of leaves or needles.

Biology: The life cycle of many rust fungi are very complex. In the early spring, infected dead aspen leaves on the

infects aspen leaves in the summer. Yet another spore, the golden-yellow uridiniospore, appears on both sides of infected aspen leaves within 2 weeks. This is called the repeating stage because these spores germinate to reinfect aspen leaves throughout the growing season. An overwintering spore is produced on the underside of infected leaves in late fall.




Figure 177. Rust pustules of Melampsora sp. on aspen leaves.

Effects: During wet summers this fungus can cause the leaves of highly susceptible clones to shrivel and drop prematurely, reducing growth.

Similar Insects and Diseases: Willows are host of several species of *Melampsora*, each which has a different alternate host. These species cannot be differentiated morphologically on the willow.

References: 38, 92

Shepherd's Crook Venturia spp.

Host: Aspen

Symptoms/signs: Curling, blackening and dieback of the tips of terminal and lateral shoots on young aspen sprouts.

Biology: Primary infections by *Venturia* spp. occur in spring on leaf blades or petioles and on young stem tissue. Dark brown to black lesions expand rapidly, causing leaves and shoots to droop, whither, and become brittle. Secondary infections, promoted by wet weather, are initiated throughout the period of shoot elongation by conidia from newly blighted shoots. New shoots frequently grow adjacent to blighted ones and are blighted in turn. Lesions do not extend into woody twigs.

Effects: In wet seasons, *Venturia* spp. can kill virtually all terminal shoots in susceptible aspen stands that are regenerating by sprouting. This damage reduces height growth and deforms trees

by causing a bend in the stem at the point where a lateral shoot became a new leader following death of the terminal shoot. Successive leaders may be killed during seasons of severe disease. Plants less than 3 meters tall are at greatest risk, and damage becomes negligible as trees attain heights greater than 5 meters.

Similar Insects and

Diseases: This is the only fungus on aspen causing black lesions on leaves, petioles and stems, which curl and form characteristic shepherd's crooks.

References: 38, 92

Figure 178. Curling and blackening of aspen shoots from Shepherd's crook.

Sycamore Anthracnose Apiognomonia veneta (Sacc. & Speg.) Höhn

Hosts: Arizona

sycamore

Symptoms/signs:

Sycamore anthracnose has a range of symptoms corresponding to the three phases of this disease. The symptoms include: cankers on buds and twigs; shoot blight following a period of cold spring weather; and leaf blight from direct infection of leaves. Foliar lesions characteristically



Figure 179. Leaf blight caused by sycamore anthracnose.

extend along the veins and involve interveinal tissue. Sometimes large irregular marginal lesions develop.

Biology: *Apiognomonia veneta* enters twigs via petioles during the growing season and remain quiescent until host dormancy, when it colonizes and kills bark and cambium. In the early spring,

Figure 180. Chronic infection by sycamore anthracnose results in repeated dieback of shoots.

spores produced in these cankers infect new shoots and leaves. Shoot blight, the second phase of the disease, involves the rapid death of expanding shoots and leaves. This phase tends to develop suddenly during a period of cold spring weather.

sycamore anthracnose is leaf blight, which results from direct infection of leaves. It starts out most severe on low branches.

weeks of growth. Hyphae of the fungus often grow down petioles into the twigs, setting the stage for the next year's damage. Effects: Repeated twig dieback alters the form of sycamores in two ways. First, when the terminal twig on a branch is killed and a lateral takes over as the new leader. Since infection takes place repeatedly during the life of a susceptible tree, the branch axis changes direction again and again, and crooked branches result. The second altered form is the development of a cluster of twigs around a common point on a branch because of the repeated killing of terminals. **Similar Insects** Figure 181. Severe defoliation by and Diseases: Sycamore sycamore anthracnose. anthracnose may occasionally be confused with injury by late spring frost. However, frost damage may affect several species in the same area and Apiognomonia veneta affects only sycamore. References: 92

Figure 182. Infected trees refoliate but branch dieback is evident.

Conifer Needle Diseases

Although occasional outbreaks appear quite dramatic, long-term damage from conifer needle diseases in the Southwestern Region is usually minor. Sporulation, spread, and infection by these

pathogen, and successful infection depends on favorable weather conditions. Needle diseases are often associated with plantations, such as Christmas tree plantations, where genetic variability to resistance is limited.




Figure 183. Conifer needle diseases are most noticeable before bud break.

Lophodermella Needle Cast Lophodermella cerina (Darker) Darker

Host: Ponderosa pine

Symptoms/Signs: *Lophodermella cerina* infects the needles of seedlings, saplings, and the lower crown of larger trees. Current year needles are infected and turn brown and die by early summer of the following year. Following bud break, green current year needles and brown previous year needles give the tree a distinctive appearance. The brown needles are "cast" prematurely compared to normal needle drop. Successive years of infection occur only with favorable weather conditions and results in thin crowned trees.

Biology: Fruiting bodies are short, oval, light brown, and easily overlooked. The spores mature after the needles have been cast. They infect new needles shortly after bud break. Spores germinate and infect needles directly through the epidermis. Infected needles extend normally and are usually killed by the fungus before the next year's needles emerge.

Effects: Although rare, growth loss is the primary long-term effect of needle cast. Successive years of infection are usually the result of repeated abnormally high rainfall in spring and early summer.

Figure 184. Lophodermella needle cast on 1-year needles before bud break.

Though trees can be killed if infection is repeated for several years in a row, they are more often predisposed to other pests.

Similar Insects and Diseases: From a distance, trees with needle cast symptoms may be mistaken for those affected by winter damage, salt damage, or needle miners. On close inspection, needle cast needles are completely brown with dark fruiting bodies present. Winter damage, salt damage, and needle miner needles often have brown needle tips with green bases.

References: 71, 82, 92




Figure 185. Diseased 1-year needles are cast in late summer leaving current year and 2-year+ needles.

Elytroderma Needle Cast Elytroderma deformans (Weir) Darker

Hosts:

Symptoms/Signs:

Effects: Although elytroderma needle cast is known as an occasional disease of ponderosa pine in the Southwest, it is common in some areas. In mature trees, if less than one-fourth of the branches are infected there is little or no effect on the tree, but if more branches are infected, needle cast alone can kill a tree or predispose it to attack by bark beetles.

Similar Insects and Diseases: The witches' brooms produced by elytroderma needle cast are similar to those caused by dwarf mistletoe, however, the existence of dead second year needles are indicative of needle cast and not dwarf mistletoe infection. Elytroderma is the only needle cast that is perennial.




Figure 187. Cambium of infected branch has pockets of dark resin.

Red Band Needle Blight Mycosphaerella pini Rostr. in Munk (Scirrhia pini Funk & A.K. Parker) Dothistroma septospora (Doroguine) Morelet (D. pini Hulbary)

Hosts:

Symptoms/Signs:

bands around the needles, which later turn red. Infection of current and second year needles is usually in the lower crown of sapling size trees. Seedlings and large trees are rarely infected. Newly infected needles have green bands that turn red or brown in late summer. These needles die from the tips back. Infected second and older year needles can be cast the same year they become infected.

Biology: Needles are infected during rainy periods by rainsplashed spores. The number of infection cycles depends on the climate. During wet years, several cycles of infection can occur. The *Dothistroma septospora* stage is the asexual, imperfect stage,

and the one most commonly found.

Effects: Dothistroma needle cast has been observed on

in New Mexico. It is most often associated with offsite planting. Several years of severe infection results in reduced growth and death of infected trees.

Similar Insects and

Diseases: Infected trees can look like the lower crown was sprayed with herbicides. The death of needles from the tips to fascicle is similar to the affects caused by high concentrations of de-icing salts.

References: 77, 82, 92

Figure 188. Dothistroma blight of Austrian pine.

Rhabdocline Needle Cast Rhabdocline spp.

Host:

Symptoms/signs: Tiny chlorotic spots on one or both

summer or early autumn. These lesions enlarge and darken to purplish brown spots and bands that are conspicuous by late autumn to early spring. Numerous lesions coalesce and involve the entire needle except for a short basal portion. Discolored foliage is most conspicuous the following spring.

Biology: Rhabdocline species produce apothecia May through June in needles still attached to the twigs. The apothecia appear as swellings up to several millimeters long and open by splitting the needle surface, usually near the midrib. Ascospores are airborne and penetrate developing needles directly through the cuticle. Only one infection period occurs per year, but it may




Figure 189. Rhabdocline needle cast lesions on 1-year Douglas-fir needles.

last several weeks. Needle colonization increases during the fall as symptoms become apparent. Infected needles drop during the winter, making missing foliage the only noticeable sign of disease.

Effects: This is the most important needle disease of

and ornamental nursery stock. Seedlings infected in nurseries can become foci of subsequent damage in plantations. Occasional outbreaks in forest stands can appear spectacular, but trees usually recover.

Similar Insects and Diseases: The fruiting body of Rhabdocline may be confused with secondary fungi.

References: 82, 92

Figure 190. Needle cast due to rhabdocline needle cast.

True Fir Needle Cast Lirula abietis-concoloris (Mayor ex Dearn.) Darker

Host:

Symptoms/signs: Dark brown or black elongate fruiting bodies are found on second year needles. The fruiting bodies extend down the center of the lower needle surface for almost the full length.

Biology: Infection begins on young, developing needles during periods of rainfall. Fruiting structpn3aauctpns ot tsees

White Pine Needle Cast Lophodermella arcuata (Darker) Darker (Hyphodermella arcuata Darker)

Host: Southwestern white pine

Symptoms/signs:

brown to concolorous, elongate to elliptical, fruiting structures,

which develop on previous year's dead and dying needles.

Biology: Fruiting bodies in dead, second year needles open by means of a

warm wet weather and release colorless spores that have sticky, gelatinous sheaths. Airborne spores settle and penetrate young, developing needles. The following spring, diseased needles turn reddish brown and by July, they are straw colored.

Effects: Lophodermella arcuata infects only the current year's needles, and a single attack results in only partial defoliation of the host. Repeated consecutive infections are rare, but can lead to reduced tree growth and vigor.

Similar Insects and

Diseases: This is the only known needle cast disease of white pine in the Southwest.

References: 82, 92




Figure 192. Lophodermella arcuata cause browning of 2-year needles.




Figure 193. Black fruiting bodies of Lophodermella arcuata.

Pinyon Needle Rust Coleosporium ribicola Arthur (C. jonesii (Peck) Arth.)

Hosts:

alternate hosts.

Symptoms/Signs:

On ribes, uredinia appear as small, round, golden yellow pustules during the summer. Telia, which are waxy and darker orange,

develop in late summer on the same ribes leaves.

Biology:

spore stages on two different hosts to complete its one year life cycle. Spores from ribes leaves infect pine needles in late summer to early fall, where the fungus overwinters. Orange

Figure 194. Aecia of Coleosporium ribicola on piñon.

droplets form on infected needles in early spring and bright orange aeciospores by late spring. The latter are wind disseminated and infect newly emerged ribes leaves. Urediniospores develop on the ribes and multiply during wet periods. Telia develop in late summer

Effects: Affected needles drop prematurely. This disease *Ribes*

spp.

Similar Diseases: *Coleosporium ribicola* on ribes can be mistaken for other rusts of ribes, such as white pine blister rust, *Cronartium ribicola*

region, *Coleosporium crowellii*, which is microcyclic, meaning there is no alternate host. The fruiting structures are more reddish-orange and waxy rather than powdery as in *Coleosporium ribicola*.




Figure 195. Uredinia and telia of Coleosporium ribicola on ribes.

ecay of wood is caused by fungi that utilize the cell wall components of xylem as a source of

break down tissues in branches, roots, and stems of trees. They play an irreplaceable role in the earth's carbon cycle by returning to the atmosphere billions of tons of carbon each year and provide valuable habitat for many species of wildlife. Although vital to ecosystem functioning, decay can render trees useless for wood products and create hazard trees in recreation sites.

Many decay fungi discussed here decay the heartwood of living trees, others decay the wood of dead trees, and some grow on both live and dead trees. Most decay fungi do not interfere with the




Figure 196. Evidence of decay is marked by fungal conks or cavity nesting holes.

normal growth of live trees since they stay in the heart or xylem wood of the host. However, they can affect tree structure and contribute to wind breakage and windthrow as decay progresses. There are some decay fungi that cause mortality. These fungi begin decaying the host as a heart rot and later move into the sapwood and

tree. Other fungi decay the wood of dead trees and can also decay dead sections of live trees, such as a dead top or old wound on the bole.

Decay levels in live trees are highly correlated with age. For example, high amounts of decay are typically found in sites where conifer trees are 150+ years old. Susceptibility to stem decay fungi varies with tree species. Some hardwoods—like cottonwood and aspen—are the most susceptible to stem decay in the Southwest, and

ponderosa or southwestern white pine. However, susceptibility is also dependent on site quality and host genetics; silviculturists have observed some 70-year-old ponderosa pine sites with high levels of decay.

Red Ring Rot Phellinus pini (Thore.:Fr.) A. Ames

Hosts:

Symptoms/signs: *Phellinus pini* produces swollen knots where branches were previously shed, irregular bulges with exuding

bodies at branch bases, branch stubs, knots, wounds, and cracks. These fruiting bodies, or conks, are bracket-like with a brownish-black upper surface, concentric furrowed rings and a brown

western larch, pines, and some spruces. A punk knot is a mass of tightly packed sterile brown hyphae that extends from a decayed branch stub within the trunk to a local swelling on the surface.

Biology: Decay often begins near the junction of heartwood and sapwood and may extend into sapwood adjacent to wounds. The

one central column or in several discrete columns that extend from branch stubs.



Figure 197. Phellinus pini on Engelman spruce.

Effects: *Phellinus pini* is thought to be the most common trunk decay fungus of conifers in North America. Decay columns commonly extend 10 meters or more, rendering entire trunks useless for lumber. Decay is much more extensive in old trees in virgin and unmanaged forests than it is in managed forests where trees are harvested at relatively young ages.

References: 27, 30, 92

Figure 198. Red ring rot on Douglas-fir.

Red Rot Dichomitus squalens (Karst.) Reid. (Polyporus anceps Pk.)

Hosts:

Symptoms/signs:

fruiting body on the underside of dead branches or stems with intact bark. The pore surface is white when fresh and ages to yellow. The

red rot fungus causes a white pocket rot. Like other wood decays, it has two distinct stages: incipient and advanced.

The incipient stage is characterized by a reddish-brown discoloration of the affected wood, unaccompanied by any obvious changes in structure or strength. The advanced stage is characterized by




Figure 199. Red rot fruiting body on the underside of a dead ponderosa pine branch.

accompanied by progressive changes in structure and reduction in strength. As decay progresses, the pockets become more and more numerous until they merge and give the affected wood the

material disappears, leaving the bleached, grayish-brown, decayed wood in either a stringy or a somewhat amorphous condition.

Both stages of red rot are usually visible in a board sawed from a decayed log. At the point where rot started in the trunk heartwood, advanced decay often forms a cavity. Extending in both directions from this point are more or less continuous columns of advanced decay, bordered by incipient decay.

Biology: The spores are dispersed by wind, land in cracked bark crevices of dead branches, and germinate to colonize the area between the bark and wood and eventually the dead wood, provided it has intact bark. The red rot fungus fruits abundantly on the lower side of decaying dead material in close contact with the ground. The

then develop annually during the rainy season for about 6 years.

Effects: Dichomitus squalens is the most common decay of ponderosa pine in the Southwest. It is a decayer of slash (a saprophyte) as well as a heartrot in live trees. It has been reported to cause a

in live trees 150+-years-old when grown for timber. As with other decay fungi, *D. squalens* provides habitat for cavity nesting birds and other wildlife. Although this fungus is of minor occurrence on living trees in second growth forests, there




Figure 200. Red rot decay is typically associated with old growth ponderosa pine.

are reports of extensive decay of young trees (<80 years) in localized areas.

Similar Insects and Diseases: Many fungi decay ponderosa pine slash, but *D. squalens* is probably the most common.

References: 27, 29, 51, 92

Indian Paint Fungus Echinodontium tinctorium (Ell. & Ev.) Ell. & Ev.

Hosts:

Symptoms/signs: *Echinodontium tinctorium* produces woody, perennial hoof-shaped conks, 4-20 cm in diameter, usually at the undersides of branch stubs. These fruiting structures sometimes form at wounds or on limbs. The upper surface is dark gray-to-black, rough and cracked. The lower surface is gray-brown to black, composed of thick, blunt, tooth-like spines. The interior tissue is brick red to rust red, from a pigment that extends into the adjacent decayed wood. The common name for *Echinodontium tinctorium*, "Indian paint fungus," is derived from the Native American's use of conks in the preparation of red paint pigments.

Biology: The spores of *Echinodontium tinctorium* are dispersed during cool, wet weather in the fall. They remain viable during winter, germinating best after a period of freezing temperatures. The fungus enters branchlets between 0.5 mm to 1.5

mm in size and spreads through the pith to the main branch. Here the fungus can remain in a dormant condition for 20, 50, or even 100 vears. When the main branch dies or breaks off.




Figure 201. Echinodontium tinctorium conk showing orange inner-tissue.

the fungus becomes active again, and decay of the heartwood portion of the main branch begins, eventually spreading to the main stem of the tree.

Effects: This fungus is the main cause of heart rot and

defect and are associated with substantial volumes of decay. One fruiting body usually indicates that the entire cross section of the log is decayed for a distance of 2 m above and 2.5 m below the conk. Decay may also be present in trees that do not bear sporophores. The rot is most common in the mid-trunk but may also extend into the butt or down from the top. In very late stages of decay, the trunk may become completely hollow. Infected trees that break or are windthrown are important sources of hollow logs for wildlife habitat and act as nurse logs for regeneration.

Similar Insects and Diseases: Advanced stages of decay closely resemble equivalent stages of rot associated with *Stereum sanguinolentum*. However, *S. sanguinolentum* fruiting bodies are completely different, as they form thin, crust-like layers with a grey to light brown gilled lower surface that turns blood red when bruised.

References: 21, 29, 92, 100

Red Belt Fungus Fomitopsis pinicola (Swartz:Fr.) Karst. (Polyporus pinicola Swartz:Fr.)

Hosts: Many conifers and aspen

Symptoms/signs: *F. pinicola* forms perennial conks that are

near the white to cream colored edge. These fruiting bodies form on dead trees and logging slash and not on infected living trees. The decay is a brown cubical rot with shrinkage cracks in which prominent white sheets of mycelium develop.

Biology: This fungus enters living trees with airborne spores colonizing and infecting through wounds and broken tops. It may be introduced into dying or dead trees by insects, as it has been isolated

galleries.

Effects: F. pinicola is one of the most important brown rot pathogens of old-growth western conifers, but it acts slowly and is not considered a major decay pathogen of secondgrowth forests.

However, it is an important component

Figure 202. Fomitopsis pinicola on dead Douglas-fir.

of the coniferous forest ecosystem because it decays dead trees and logging slash and leaves a lignin-rich residue that is very stable and

in the upper layers of soil. This residue enhances water holding and cation exchange capacities of soil and is a favorable habitat for the

Similar Insects and Diseases: Brown rot caused by *F. pinicola Phaeolus schweinitizii P. schweinitzii* is a root and butt rot pathogen, it is found in the lower 3 meters of the trunk and in the roots, and it fruits from the roots out through the soil of live trees. *F. pinicola* only fruits on dead wood material so is frequently observed on stumps.

References: 27, 29, 92

Pouch Fungus Cryptoporus volvatus (Peck) Shear

Hosts: Ponderosa pine and other conifers

Symptoms/signs: *C. volvatus* produces small, creamcolored to tan, leathery fruiting bodies. The most distinguishing feature of *C. volvatus* is that it has a fungus sheath with a single hole covering the lower spore-bearing surface. This is not a heart-rot fungus, but causes a grayish white rot of the sapwood of recently killed trees.




Figure 203. Cryptoporus volvatus on dead ponderosa pine.

Biology:

Insects enter, feed, and then exit fruiting bodies of the pouch fungus, carrying spores to infect recently killed or dying trees. Typically, it occurs only on dead trees and snags within 1 to 2 years after the tree's death.

Effects:

Most commonly found on ponderosa pine in the Southwest, but can infect other conifers. This is not an aggressive pathogen, but invades

in trees killed by bark beetles.

Similar Insects and Diseases: No other conifer pouch

References: 7, 27, 29

Blue Stain Fungi Ophiostoma spp. Ceratocystis spp.

Hosts: Conifers

Symptoms/signs: Blue-green discoloration of sapwood (often in wedge shapes) in recently killed trees. Blue stain fungi frequently originate from bark beetle galleries.

Biology: Blue stain fungi are carried by bark beetles and other wood inhabiting insects and are associated with tree mortality. They are mostly blue-staining and primarily from the genera *Ophiostoma* and *Ceratocystis*. The spores germinate and produce a mycelium (thread-like mass) that colonizes the phloem and sapwood, eventually blocking the water-conducting columns of the tree. The formation of a wedge-shaped stain is due to movement of hyphae along the rays from the outside of a log.

Effects: Stain fungi often hasten the death of trees attacked by bark beetles. The stain may result in a reduction in the value of timber or timber products by discoloring sapwood, but does not affect wood strength.

Similar Insects and Diseases: Another group of sapstaining fungi are wind disseminated, of various colors (blue, brown, or gray) and are from the genera *Aureobasidium* and




Figure 204. Blue stain fungi invade the sapwood.

Alternaria. Black stain fungi, which are closely related to blue stain fungi, also block water-conducting columns. However, black stain fungi spread to healthy trees by root-to-root contact.

References: 92, 100

False Tinder Conk Phellinus tremulae (Bondartsev) Bondartsev & Boresov in Bondartsev

Host: Aspen

Symptoms/signs: *P. tremulae* produces woody conks on aspen trunks, typically at branch stubs or scars. The conks are hoof

shaped, brown to black, with a rough, cracked upper surface and tan to white pore surface. The conks are usually attached to the host by a granular core of tissue that continues into the decayed branch within the trunk. *P. tremulae* also produces hard, blackish sterile masses of mycelium (sterile conks) at branch scars, hence the name. The yellowish white spongy decay caused by this

central core.

Biology: Airborne spores of *P. tremulae* infect fresh branch stubs or wounds. The fungus decays the dead or injured area and then gains entry to the heartwood.

Figure 205. Phellinus tremullae conk on aspen.

Effects: *P. tremulae*

is the most common cause of aspen stem decay in the Southwest. It causes less mortality than aspen canker fungi or ganoderma root rot but can create hazard trees in recreation sites.

Similar Insects and Diseases: No other fungi with the characteristics of the fruiting body are found in aspen.

References: 30, 38, 92

Heart Rots of Oak Inonotus dryophilus (Berk.) Murr. Phellinus everhartii (Ell. & Gall.) A. Ames.

Hosts: Oaks

Symptoms/signs: Oaks are host to many heartwood decay fungi. Two of the most common are *I. dryophilus* and *P. everhartii*. The former produces an annual fruiting body that degrades quickly following spore dispersal, while the latter produces a perennial

fruiting body (conk) that grows a fresh sporulation layer every year.

Biology: As with most decay fungi, branch stubs are the most common sites of infection, but entry also occurs through trunk injuries. Both fungi produce fruiting bodies on living trees.

Effects: Both fungi are capable of causing extensive heartwood decay.

Figure 206. Inonotus dryophilus on Emory oak.

Similar Insects and Diseases: There are many other fungi that decay the heartwood of living oak trees, but these are the most common.

References: 29,

92

Figure 207. Heart rot, Phellinus everhartii, on gambell oak.

Canker Rot of Oak Inonotus andersonii (Ellis & Everh.) Cerny

Hosts: Oaks
Symptoms/signs:

I. andersonii
bright yellow to yellowbrown fruiting bodies with
peg-like outgrowths beneath
the bark or outermost layers
of wood. These fruiting
bodies degrade quickly,
turning dark dull brown to
black, as if the wood was
burned.

Biology: Branch stubs are the most common sites of infection, but entry also occurs through trunk injuries. *I. andersonii*

causing a white rot, and then moves outward, killing the sapwood and cambium (hence the name, canker rot). Trees often break at these cankers. Figure 208. Inonotus andersonii forms bright orange fruiting structures beneath the bark of infected trees that blacken with age.

Effects: *I. andersonii* is a common decay pathogen of oaks throughout the United States, but also causes mortality where decay continues out into the cambium, girdling infected trees.

Similar Insects and Diseases: Old fruiting bodies of *I. andersonii*

References: 29, 92




Figure 209. Canker development in aspen begins with discoloration of the bark.

Black Canker Ceratocystis fimbriata Ellis & Halst.

Host: Aspen

Symptoms/signs: Cankers are target shaped with concentric ridges of dead callus. Older cankers typically have a central area of dead wood surrounded by a series of bark calluses. These callused areas may be concentric in outline, but usually are irregularly shaped

and ragged in appearance

because of the massive callus

Bark and exposed wood darken, resulting in a black canker.

Biology:

Ceratocystis fimbriata can infect through the epidermis of leaf blades, petioles, and young stems; but trunk wounds are considered to be the primary places of infection. Several species of beetles have been found to carry the fungus and are believed to vector it.

a circular necrotic area on the trunk around a fresh wound or branch junction. During cambial growth in

Figure 210. Callus folds and flaring


dead bark caused by Ceratocystis

the spring, the tree forms a callus at the margins of the canker, which temporarily walls off infection. The fungus invades the new cambium and inner bark during the tree's next dormant season and kills a new zone of tissue. This process is repeated each year until

the canker, consisting of successive rings of dead bark and wood, is formed. The fruiting structures, perithecia, are formed in the

spring along the border of the canker on tissues dead at least a year. Spores ooze from the perithecia in sticky masses and are often vectored to other wounded trees by insects. Boring insects are often found in cankered areas. There appears to be a genetic predisposition to infection, as some aspen clones are more susceptible to infection than others.

Effects: *C. fimbriata* seldom kills large trees since trees generally grow faster in circumference than black cankers enlarge. However, it is possible for several cankers to coalesce and girdle a tree. The greatest damage from this disease is stem deformation and subsequent decay that can result in bole breakage.



expose target canker.

Similar Insects and Diseases: Sooty bark (*E. pruinosa*) is also black in color but does not form a target canker like Ceratocystis.

References: 38, 39, 44, 92

Sooty Bark Canker Encoelia pruinosa (Ellis & Everh.) Torkelson & Eckblad (Cenagium singulare (Rehm) R.W. Davidson & Cash)

Host: Aspen

Symptoms/signs: Young cankers appear as slightly sunken areas with near normal bark color. Dead bark eventually falls off and reveals the black, sooty and crumbly inner bark diagnostic for this canker. Beneath the inner bark is a distinctive, black, feather-like pattern on the wood surface. The perennial cankers form a band of light gray bark that remains attached at the margin of each annual zone of expansion giving the tree a barber pole appearance. Gray to black leathery cup-shaped fruiting bodies develop on the inner bark along the canker margin. The wood beneath cankers is light gray.

Figure 212. Barber pole appearance of aspen bole infected with Encoelia pruinosa.

fungus infects trees through wounds and invades the inner bark and cambium. Cankers develop rapidly, extending as much as 1 m in length and 0.3 m in width in a year, however, the mean annual extension is about 45 cm vertically and 16 cm horizontally. The cup-shaped fruiting bodies-apotheciadevelop on the surface of old dead inner bark and open when wet. Spores are forcibly ejected and wind disseminated when moisture and temperature conditions are favorable. These spores can cause new infections.

Biology: The

Effects: *Encoelia pruinosa* is considered the most lethal and aggressive pathogen of aspen in the West because it can kill trees of all sizes in 3 to 10 years; some mature trees have been observed to die in 4 to 5 years. Sooty bark is found mainly on the larger dominant and codominant trees older than 60 years, in the middle elevational limits of aspen. Cankers are more common

Figure 213. Cup-shaped fruiting bodies of sooty bark canker.

camper damage, or animal damage.

in stands

disturbed by

partial cutting, construction.

Similar Insects and Diseases: Cryptosphaeria canker can also produce dead inner bark with a soot-like appearance. However, Cryptosphaeria produces a long lens-shaped, light-colored area, which lacks the barber pole design.

References: 38, 39, 44, 92

Cytospora Canker Cytospora chrysosperma (Pers.:Fr.) Fr.

Hosts: Aspen, cottonwood, alder and other riparian species

Symptoms/signs:

Young cankers on smooth bark appear as brownish-yellow sunken areas. They are often irregular in outline and range from diffuse to slightly target-shaped. Later, the bark often splits at canker margins, the inner bark turns black, and wood beneath the canker is stained brown and water soaked. Black, pimple-like, asexual fruiting bodies (pycnidia) develop within a few weeks after death of the bark. They extrude spores in orange-red tendrils in wet




Figure 214. An expanding cytospora canker.

weather. Later, black, pimple-like sexual fruiting bodies (perithecia) originate beneath the bark, which expel white masses of spores during wet weather that collect on the surface of the bark.

Biology: The fungus causing cytospora canker exists in two stages, a sexual stage known as *Valsa sordida*, and an imperfect, or asexual, stage called *Cytospora chrysosperma*. The latter is more commonly encountered. The fungus is considered a normal

parasitizes bark that has been injured or weakened by any cause. Trunk cankers are formed by a gradual killing of the bark in a more or less circular area over a period of several years. Annual canker growth can be seen by the slight annual callus formation around the

perimeter of infection. The fungus fruits readily in the dead outer bark; even when typical canker symptoms fail to develop. The *Cytospora* stage forms small black fruiting bodies from which sticky spores ooze out in long, coiled, orange-to-dark-red masses called

formed beneath and in a circle around the old pycnidia. Some ascospores are forcibly discharged from the perithecia; others collect around the openings of the perithecia in sticky white masses on the dead bark. The diseased inner bark remains attached to the tree for 2 or 3 years before turning lighter brown in color and falling off in large pieces.

Effects: Cytospora chrysosperma is the most common fungus found on aspen throughout its range. It is weakly parasitic

and normally attacks stressed trees.

Cytospora infection is associated with frost cracks and sunscald, elk feeding wounds, tree vigor and damage

Similar Insects and Diseases:

The orange tendrils or spore horns distinguish this fungus from all of the other aspen canker causing fungi.



Figure 215. Cytospora spore tendrils exuding from bark of a dead tree.

References: 38, 44, 92

Cryptosphaeria Canker Cryptosphaeria lignyota (Fr.:Fr.) Auersw.

Host: Aspen

Symptoms/signs: Young infections and canker margins are orange to light brown. The cankers are long (3 meters or more) and narrow (2-10 centimeters wide), appearing as grayish depressions in the bark, with callus ridges forming at the edge. Dead bark adheres to the canker face. Inner bark is black and sooty with obvious

mm). Light orange, tiny asexual fruiting bodies may form near the edge of the canker.

shaped sexual fruiting bodies (perithecia) develop beneath bark dead for more than 1 year. *Cryptosphaeria populina* also causes stain in the sapwood and heartwood and causes a yellow-brown, mottled decay.

Biology: Spores of *Cryptosphaeria* populina are released during wet weather and infect fresh wounds in the inner bark and wood. The fungus eventually colonizes sapwood and

Figure 216. Cryptosphaeria colonizes sapwood and cambium.

heartwood, causing discoloration and decay, before penetrating the bark and causing a canker. Brown, mottled decay develops in the central part of the column of discolored wood. Mortality occurs not from bark necrosis as it may seem, but because the pathogen kills a large volume of sapwood.

Effects: Trees up to 15 centimeters d.b.h. may be killed within 3 years, with older trees taking longer to girdle. Branch cankers are often found on large trees, where they girdle the branch and enlarge onto the trunk. The decay associated with the canker predisposes infected trees to wind breakage.

Similar Insects and Diseases: Bark that has been dead for more than 1 year from *C. lignyota* is black, stringy, and sooty-like, similar to sooty-bark canker (*E. pruinosa*). However, they are easy to distinguish, because of the lenticular-shaped fruiting structures and barber pole design of *E. pruinosa*, both lacking in a Cryptosphaeria canker.

References: 38, 44, 92

Hypoxylon Canker Hypoxylon mamatum (Wahlenberg) J.H. Miller

Host: Aspen

Symptoms/signs: Young infections occur as slightly sunken, irregular, yellowish-orange areas around wounds or branch stubs. The underside of diseased bark appears laminated or mottled black and yellowish white. White mycelial fans form under the bark and are most evident at the top and bottom edges of the canker. Gray fungal pillars (asexual fruiting bodies) form beneath loose, blistered bark. Later, patches of gray-black, crust-like mounds form on the wood. These mounds contain perithecia, sexual fruiting bodies. The papery outer bark sloughs from older infections, exposing a blackened, crumbly inner bark. The cortex in the central portion of older cankers cracks in a checkerboard fashion, sloughing off in small patches.




Figure 217. A section of a Hypoxylon canker with cracked and blistering bark.

Biology: In the humid Lake States area, the fungus invades the sapwood and trees often die before they are completely girdled because sapwood decay beneath a trunk canker predisposes the tree to wind breakage. In the arid Southwest, however, breakage is not common. Cankers on large trees in the Southwest may attain ages of 20 to 50 years before tree death. A live 1 m d.b.h. aspen tree in Arizona was observed with a Hypoxylon canker extending from the ground to a height of about 12 m, only half-girdling the tree. There appears to be a genetic predisposition to infection, because some clones are more susceptible to infection than others within the same geographic area.

Effects: Hypoxylon canker disease is the most important canker disease of aspen in the Lake States region, but of minor importance in the Southwest. Hypoxylon canker annually kills an estimated 1 to 2 percent of the standing aspen volume in the Lake States area. While the disease causes serious mortality in localized areas in the Southwest, mortality rates have not been determined.

Similar Insects and Diseases: It is the checkerboard pattern of older infections that distinguishes this canker disease from others that occur on aspen.

References: 38, 44, 92

Atropellis Canker Atropellis piniphila (Weir) Lohman and Cash

Hosts: Southwestern white pine, limber pine and ponderosa pine **Symptoms/signs:** Cankers appear as elongated depressions covered with bark. Atropellis species produce two kinds of fruit across, and contain multiple chambers where spores form. The cupshaped structures are black with a brown interior, measuring 2-4 mm across when moist. They form on cankers 2 or more years old and are present year round. Dark brown mycelium permeates the bark and wood, creating a zone of bluish black stain that in cross-section is initially wedge shaped, pointing toward the center of the stem,

Figure 218. Fruiting structures on atropellis canker.

becoming irregular with age. Species are separated based on spore morphology.

Biology: Infection begins in a branch axil on stems 5-30 years old and so there is typically a branch stub in the center of the

the fungus to infect wounds or bruised bark as well as apparently undamaged bark of young stems.

Effects: Atropellis canker can be found throughout the region but its occurrence is infrequent in natural forest settings. Although Atropellis rarely kills trees, there are areas where outbreak levels are reached and mortality occurs in young trees. It can be damaging in Christmas tree plantations where it causes trunk deformities and economic loss.

Similar Insects and Diseases: Atropellis canker may be confused with other canker-causing fungi; however, atropellis is the only one that stains the wood.

Reference: 100

here are several species of stem rust fungi on conifers in the Southwest. The rust fungi are unique in many respects. Some species of rust fungi are autoecious, which means they complete their life cycle on one host. Other species are heteroecious, which means they complete their life cycles on two entirely unrelated hosts. The most important rust species of trees are those that invade the main stem and cause mortality. The most destructive stem rust listed below is white pine blister rust, a nonnative pathogen

found in New Mexico.

There is one species of cone rust in the Southwest and it infects Chihuahua pines in the southeast corner of Arizona.




Figure 219. Branch flagging and top-kill are common signs of stem rust infection.

White Pine Blister Rust Cronartium ribicola J.C. Fisch.

Hosts: Southwestern white pine is the only species currently affected in the Southwest; limber pine and bristlecone pine are also susceptible. Currants and gooseberries (*Ribes* spp.) are alternate hosts.

Symptoms/Signs:

Tapered branch swellings or stem swellings (on young trees) are an early symptom on pine. The characteristic white blisters (aecia) appear on mature cankers in late spring. After the blisters disintegrate, cankers have a dark, roughened



Figure 220. Cronartium ribicola spores are discharged from sacs or blisters in the spring.

appearance. Flagging (i.e. recently killed branches with red needles) occurs several years after initial infection.

Biology: Spores produced on *Ribes* leaves are wind dispersed and infect pine needles. The fungus grows into the inner bark, forming a canker that eventually girdles the branch or stem. Blisters erupt through the bark of the canker, releasing the spores that infect *Ribes*.

Figure 221. White pine blister rust sporulating on the bole of an infected tree.

Effects: This nonnative disease is one of the most damaging tree diseases in North America. Trees of all sizes can be affected, although smaller trees are killed more rapidly than larger trees. Blister rust was found near Cloudcroft, New Mexico in 1990, and is now widespread throughout the Sacramento and adjoining White Mountains. It is most common on moist, mixedconifer sites above 2,450 m. The disease has also been

found in the nearby Capitan Mountains, on Gallinas Peak

Figure 222. White pine blister rust cankers eventually girdle infected stems and branches.

Figure 223.
Cronartium ribicola on ribes.

(near Corona, NM), and most recently on the Gila National Forest, 5 km from the Arizona border. However, it has not yet been found in northern New Mexico or in Arizona (as of 2005).

Similar Diseases: Atropellis canker, which is fairly common on white pines in the Southwest, forms slight swellings with roughened bark. It produces small black fruiting bodies near the canker margin. Dwarf mistletoe causes branch swellings that

can look remarkably similar to blister rust cankers. Sunscald and mechanical injuries can

blister rust damage. *Ribes* leaves are often infected by other rusts, such as *Coleosporium ribicola*, that can be mistaken for white pine blister rust.

References: 26, 70, 92

Figure 224. Top-kill and flagging on mature white pine.

Limb Rust Cronartium arizonicum Cummins

Host: Ponderosa pine

Symptoms/Signs: Dead and dying branches, often in the middle of the crown, are quite characteristic of this disease. Sporeproducing sacs (aecia) are produced on live infected branches, and erupt through the bark in spring to midsummer. Trunks are not affected by this disease even though the fungus travels through it to get to the branches.




Figure 225. Live limbs bearing Cronartium arizonicum spore sacs in the spring/summer die in the fall.

Figure 226. Limb rust infection typically starts mid-crown and spreads in both directions.

Disease from this pathogen results in a progressive invasion and killing of branches by the fungal mycelium that is perennial in the trunk, but does not injure it. From an initially infected branch, the parasite grows into the wood of the trunk and advances upward and downward up to 18-21 cm per year in each direction and then out into branches where it sporulates. Infected branches typically die the following year.

There are at least three varieties of limb rust in the western U.S., collectively known as *Peridermium filamentosum*. *Cronartium arizonicum*, which uses Indian paintbrush (*Castilleja* spp.) as an alternate host, is the most common variety in Arizona and New Mexico. Another variety that spreads directly from pine to pine has been detected on the Kaibab Plateau in northern Arizona.

Effects: Although the disease can be very damaging to individual trees, its occurrence in an area is usually low. In Arizona, it has been observed throughout the ponderosa pine type, from the Coronado National Forest to the North Kaibab Ranger District.

Similar Diseases: Many agents can cause branch mortality, but limb rust is distinctive because of the progressive mortality in the center of the crown.

References: 79, 92, 116

Western Gall Rust Peridermium harknessii J.P. Moore Endocronartium harknessii (J.P. Moore) Y. Hiratsuka

Host: Ponderosa pine

Symptoms/Signs: This fungus forms round or pear-shaped galls on the main stem or branches, which emit orange spores. There is a white-spored race in Arizona and New Mexico that sometimes develops elongated galls.

Biology: This disease spreads directly from pine to pine. Infection occurs on a needle-bearing twig. The fungus stimulates

the production of xylem, forming galls, which enlarge over the years. Spores appear on the galls in late spring.

Effects:

This disease is probably widespread in the region, but at very low levels. Occasionally it has been destructive to



Figure 227. White-spored gall rust displaying spore sacs and swollen branch.

seedlings and saplings, following infrequent "wave years" of new infection. Gall rust deforms but seldom kills older trees.

Similar Diseases: Dwarf mistletoe can cause trunk swellings (cankers) that are usually much longer than those produced by western gall rust.

References: 92, 116

Comandra Blister Rust Cronartium comandrae Peck

Hosts: Ponderosa pine and Mondel pine (*Pinus eldarica*) Symptoms/Signs: Symptoms of disease on pine begin with slight, spindle-shaped swellings that give rise to pustules that soon exude a mass of rust colored spores. On the nonnative Mondel/ Eldarica pine, comandra blister rust causes dieback of branches and eventual tree death. On ponderosa pine, its native host, this disease is rare but occasionally girdles seedlings and saplings. On *Comandra pallida*, the alternate native host, the rust causes pale yellow leaf and stem spots and leaf abscission.

Biology: The aecial stage develops on pine in late spring through early summer and spores disseminate by wind at that time. These spores are responsible for the primary infection of the alternate host, *C. pallida*. Rain and cool temperatures favor their germination. The next spore stage, the so-called repeating stage, occurs and builds

give rise to basidiospores in late summer, which are wind-dispersed to infect pine needles. The fungus grows into the inner bark, forming a canker that eventually girdles the branch or stem.

Effects: Although this disease causes major damage to lodgepole pine and sometimes ponderosa pine in northern Colorado

and Wyoming, it is of minor consequence to ponderosa pine in Arizona and New Mexico. However, comandra blister rust has done considerable damage to plantings of Mondel pine




Figure 228. Comandra pallida with leaf infections.

in the Prescott, Sedona and Payson areas of Arizona. Beginning in the late 1970s, Mondel pines were planted widely as a landscape tree throughout these areas, until it was realized that comandra blister rust impacted their survival. Infected *Comandra pallida* are typically observed near infected Mondel pines.




Figure 229. Comandra blister rust swollen canker.

Gymnosporangium Rusts Gymnosporangium spp.

Hosts: Junipers and Arizona cypress

Symptoms/Signs: There are eight species of Gymnosporangium rusts of juniper in the Southwest and one on cypress. Most Southwestern Gymnosporangium spp. alternate on serviceberry and/or hawthorn (Rosaceae), but one species (G. speciosum) alternates on fendlera and mock orange (Hydrangeaceae). On the evergreen host tree, these rusts cause witches' brooms, galls, other branch distortions, and dieback of twigs and branches. Brown to orange hornlike or cushionlike projections (telia) are produced in the spring; these swell and gelatinize during wet periods and are quite spectacular. On alternate hosts, the rust develops colorful spots and localized swellings on leaves, fruits, and green twigs, followed by casting or distortion and death.

Figure 230. Gymnosporangium speciosum telia on juniper.

Biology: Some *Gymnosporangium* species complete their life cycles in 1 year, but many require 2 years. Two spore stages form in spring through late summer on the alternate host. The second spore type is wind disseminated and infects juniper or cypress (depending on the rust species). *Gymnosporangium* species overwinter in their evergreen hosts, where they produce telia in

alternate host. Many of these rust species become perennial in juniper or cypress and produce telia annually on galls, swellings, or witches' brooms.

Effects: In the Southwest, these fungi generally cause minimal damage to junipers, other than some deformities. Impact on

years with adequate rainfall.

Similar Diseases: Witches' brooms caused by some of the juniper rusts could be confused with mistletoe infection. *Gymnosporangium cupressi* affects Arizona cypress in central Arizona.

References: 78, 92, 116

Fir Broom Rust Melampsorella caryophyllacearum J. Schröt

Hosts:

Spruce Broom Rust Chrysomyxa arctostaphyli Dietel

Hosts: Engelm	ann and blue spruce
Symptoms/Sig similar on their	ns: Both of these diseases appear very
respective hosts. The yellow to pale green brooms are dense and compact. Stem or branch swellings may also occur near the point of infection. Biology: Windblown spores produced on an alternate host are needed to start new infections on	Figure 231. Yellow to pale-green witches' broom are conspicuous even at the forest level.
	trees. Chickweeds (Cerastium and Stellaria spp.) are alternate hosts for while bearberry or kinnikinnick Figure 232. Expanding foliage is pale green.

(Arctostaphylos uvaursi) is the alternate host for spruce broom rust. Once a tree is infected, the fungus stimulates bud formation, leading to broom development. The brooms shed their needles in the winter and grow new ones in the spring.

Effects:

Broom rusts can be found throughout much of the Southwest on their respective hosts. They typically occur at low levels, but are abundant in some locations. Infection typically results in deformity, which is

young trees. Stem infections sometimes result in topkill and/or stem breakage.




Figure 233. Foliage yellows in late summer.





Figure 234. Needles die and drop, leaving the broom devoid of foliage during winter.

Stem and Cone Rusts of Pine

Similar Diseases: Broom rusts are sometimes mistaken for dwarf mistletoe witches' brooms. However, the former are more dense and compact, and lack mistletoe shoots. Dwarf and spruces have very limited distributions Figure 235. Chickweeds are alternate hosts for in the Southwest. fir broom rust. **References:** 92, 116

Figure 236. Topkill caused by spruce broom rust.

Stem and Cone Rusts of Pine



174

broom rust.

Stem and Cone Rusts of Pine

Southwestern Cone Rust Cronartium conigenum Hedgc. & N. Hunt

Host: Chihuahua pine and oak

Symptoms/Signs: This disease produces swollen, deformed cones on Chihuahua pine that, when ripe, are loaded with orange spores beneath a thick papery outer surface. On oak, this disease produces several different spore stages. The type most often observed has the appearance of brown to black wool on the lower leaf surface.

Biology: This fungus produces spores on swollen deformed cones (galls) of Chihuahua pine in July and August, and other spores on the undersides of leaves of Mexican blue, Dunn, Emory, gray, canyon

live, netleaf, silverleaf and Arizona white oaks throughout the summer. Cones become

of development, swell into misshapen galls of various sizes, produce no seeds and do not open. Aecia develop 2 to 3 years after infection. Galls die after aecial production but remain on Figure 239. Swollen Chihuahua pine cone infected with C. conigenum and sporulating. An uninfected cone is on the right.

the trees. Large galls usually kill the branches that bear them.

Effects: During periods of outbreak brought on by favorable weather, cone rust may kill more than 50 percent of the cones on groups of trees. This disease also affects other pine species in Mexico.

Similar Diseases: None

References: 92, 116

istletoes are parasitic plants that slowly weaken and eventually kill their host trees. There are two types of mistletoe in North America, true mistletoe (*Phoradendron*) and dwarf mistletoe (*Arceuthobium*). Both are parasitic on the stems of woody plants from which they derive water and mineral nutrients. True mistletoes produce most of their organic

nutrients by their own photosynthesis and are believed to obtain little from the host. Although photosynthetic, dwarf mistletoes are dependent on the host for carbon nutrients for growth and reproduction. Dwarf mistletoes are generally considered to be more damaging than true mistletoes; however, host mortality associated with either type of mistletoe infection can occur during periods of drought when lack of stomatal closure of leaves of the parasite causes fatal water loss of the host.

Figure 240. Male (above) and fem A. vaginatum subsp. cryptopodum	

Dwarf Mistletoes Arceuthobium spp.

Dwarf mistletoes are the most common pathogens in Southwestern coniferous forests. They are parasitic, seed-bearing plants that depend on their hosts almost completely for water and nutrients. Dwarf mistletoes are natural components of many forest ecosystems in the West, having co-evolved with their hosts for hundreds of thousands of years. There are eight species in the region, each typically having one preferred host species.

Ponderosa pine dwarf mistletoe (generally referred to as **southwestern dwarf mistletoe** to distinguish it from a different species affecting ponderosa pine in California and the Northwest; it has recently been referred to as **pineland dwarf mistletoe** by some) has long been recognized as the most damaging disease of ponderosa pine in the Southwest. It occurs in over one-third of the ponderosa

roughly one-half of the mixed conifer acreage in the region. Other species of dwarf mistletoe have more limited distributions (see table).

Hosts: See table

Symptoms/signs: Aerial shoots of dwarf mistletoe plants vary by species in size, color, and pattern of branching. For example, southwestern dwarf mistletoe of ponderosa pine are often bright

often small and inconspicuous. Host branches and stems are often swollen at the site of dwarf mistletoe infections. Witches' brooms develop from either systemic infections or as a result of discrete, localized infections. The size and extent of brooms varies among dwarf mistletoes and their host.


Biology: Dwarf mistletoes have separate male and female plants. Seeds are produced annually on female plants. These are explosively released (typically 1 to 12 meters), and stick to host material. Upon germination, dwarf mistletoes produce an endophytic

system, a specialized rootlike structure that is in contact with the phloem and xylem of host trees, from which the parasite obtains nutrients and water. Aerial shoots appear 3 to 5 or more years after

infection; the period before shoots are visible is known

Spread of dwarf mistletoe occurs both from tree to tree and within the

as the latent period.



Extensive dwarf mistletoe infection greatly reduces forest productivity. On the other hand, infection

for wildlife. Large witches' brooms can serve as ideal nesting platforms for birds and small mammals, and snags create habitat for cavity nesting birds. A few species (most notably the blue grouse) are known to eat dwarf mistletoe shoots, although none depend on it as a primary food source.

Similar
Insects and Diseases:
Some fungi cause the
formation of witches'
brooms: Broom rusts




Figure 242. Spruce dwarf mistletoe.

Figure 243. Apache dwarf mistletoe on southwestern white pine (above).

Where brooms are observed, branches should be checked for the presence of aerial dwarf mistletoe shoots to distinguish broom symptoms caused by other pathogens or physiological disorders.


References: 12, 33, 34, 76, 101




180

Principal Hosts and Distribution of Dwarf Mistletoes in Arizona and New Mexico

Common Name	Species	Principal Host	Distribution
Southwestern dwarf mistletoe	Arceuthobium vaginatum subsp. cryptopodum (Engelmann) Hawksworth & Weins	Ponderosa pine	Throughout host type
dwarf mistletoe	Arceuthobium douglasii Engelmann		Throughout host type
Western spruce mistletoe	Arceuthobium microcarpum (Engelmann) Hawksworth & Weins	Engelmann spruce, Blue spruce, and Bristlecone pine	Limited portions of host type, although common in eastern and central Arizona
Pinyon dwarf mistletoe	Arceuthobium divaricatum Engelmann		Throughout the Southwest, except southeast Arizona
Apache dwarf mistletoe	Arceuthobium apachecum Hawksworth & Weins	Southwestern white pine	Eastern and southeastern Arizona, central and southern New Mexico
mistletoe	Arceuthobium abietinum f. sp. concoloris Engelmann ex Munz		North rim of Grand Canyon National Park, Santa Catalina and Chiricahua Mts., AZ
Chihuahua pine dwarf mistletoe	Arceuthobium gillii Hawksworth & Weins	Chihuahua pine	Mostly southeast Arizona, but some in central
Blumer's dwarf mistletoe	Arceuthobium blumeri A. Nelson	Southwestern white pine	Huachuca Mts., Arizona





True Mistletoes Phoradendron spp.

True mistletoes are commonly known as the Christmas or leafy mistletoes. There are eight species in the Southwest, all within the genus *Phoradendron*. Three species occur on hardwoods,

Phoradendron macrophyllum has a very broad host range, occurring on most riparian tree species, while the other true mistletoes are genus



Figure 253. Phoradendron coryae in December.

Hosts: See table **Symptoms/signs:** True

often round in form and up to 1 meter in diameter, depending on the species. Hardwood true mistletoes have thick green leaves that are nearly oval in shape, contrasting with conifer true mistletoes, which





Figure 254. Phoradendron juniperinum with pink ripe berries.

berries are white, pink or red and are ripe from October to January, again depending on the species. Evergreen clumps of mistletoe are readily observed on deciduous trees in winter when leaves are off the trees.

Biology: Fruiteating birds distribute the seeds in their droppings or by wiping their beaks. Some

bird species swallow the fruit whole and disperse the seeds to another tree, while other bird species pick out the seed, leaving it on the host plant, and swallow only the pulp.



through which water and nutrients pass from the host to the mistletoe. It takes approximately 2 to 3 years for shoots to develop, following initial infection, and another year before the plant is producing berries.




Figure 256. Phoradendron macrophyllum on Arizona sycamore.

Effects: Young or small trees are seldom infected by true mistletoe. In nearly all cases, initial infection occurs on larger or older trees because birds prefer to perch in the tops of taller trees. Severe buildup of mistletoe often occurs within an infected tree because birds are attracted to and may spend prolonged periods feeding on the mistletoe

berries. True mistletoes are not aggressive pathogens. They use the host xylem as a water source and do not cause mortality until water availability to the host is limited. In some hosts, infected portions of

the tree often exhibit galls on branches or burls in the trunk. On oaks and cottonwoods, branch dieback is associated with galls formed by the corresponding mistletoe.

Similar Insects and Diseases: Deformities caused by canker and rust fungi can resemble those caused by mistletoe.

References: 25, 35, 84, 92

Figure 257. Phoradendron californicum has red berries that ripen in winter. Figure 258. Desert mistletoe plants are red when full of fruit. Figure 259. True mistletoes are easy to spot on deciduous trees like this mesquite tree.

Principal Hosts and Distribution of True Mistletoes in Arizona and New Mexico

Common Name	Species Name	Principal Host	Distribution
Bigleaf mistletoe	Phoradendron macrophyllum (Engelm.) Cockerell	Most riparian hardwood species, except oaks	Throughout lower elevation riparian areas of both states
Southwestern oak mistletoe	Phoradendron coryae Trel.	Oak species	Throughout live oak woodlands and lower elevation gambel oak areas of both states
Desert mistletoe	Phoradendron californicum Nutt.	Leguminous trees and shrubs (e.g. Mesquite (<i>Prosopis</i> spp.), Acacia (<i>Acacia</i> spp.), Palo Verde (<i>Cercidium</i> spp.), and ironwood (<i>Olneya</i> spp.)	Throughout the ranges of host types in Arizona and extreme southwest corner of New Mexico

Figure 260.
Phoradendron densum on Arizona cypress.

Juniper mistletoe	Phoradendron juniperinum Engelmann	All juniper species	Throughout juniper woodlands of Arizona and New Mexico
Hairy juniper mistletoe	Phoradendron capitellatum Torr. ex Tel.	Utah, alligator and red-berry juniper	Southern Arizona and southwest New Mexico
Texas juniper mistletoe	Phoradendron hawksworthii Wiens & CG Shaw	Alligator and one- seed juniper	Southeastern New Mexico
Dense mistletoe	Phoradendron densum Torr. ex Tel.	Arizona cypress	Central Arizona
mistletoe	Phoradendron pauciflorum Torr.		Santa Catalina Mountains of southeast Arizona

	Figure 261. Phoradendron hawksworthii on juniper in New Mexico.
Figure 262. Phoradendron often kills tops of infected white fir.	

Root disease fungi affect all tree species in the spruce, and aspen being the most susceptible to damage. Susceptibility also varies by tree age, genetics, type of root disease pathogen present and site history.

Root disease or decay fungi spread from roots of diseased trees to those of healthy ones. They start in a tree or stump and spread slowly outward in all directions, resulting in a slowly enlarging group of dying and dead trees. The oldest kills are located at the center of infection with a fringe of recently killed and dying

trees around the outer edge. In the Southwest, these "centers" of disease are small in size, typically less than 1 acre.

Root disease is referred to as "a disease of the site" because the fungi colonize dead and dying trees and remain in dead roots and soils for many years. Many of these fungi have the ability to act as both pathogens in



Figure 263. Fruiting bodies of root disease fungi are located at the base or roots of infected trees.

live trees and saprophytes in dead wood material. Not only are the current trees affected, but also trees that become established in the future. Some root disease fungi have been shown to remain alive and active in infested sites for more than 50 years.

The aboveground symptoms of trees affected by root disease include chlorosis, reduced needle length, progressive thinning of foliage, fading crown, reduced tree growth, and death. These symptoms are similar to those caused by drought, high water table, and bark beetle attack. However, the decline of trees affected

by root disease usually extends over a per	nod of a few to several
years and not all trees succumb at the san	ne time. The other causes
of decline kill trees more	
rapidly, generally in 1 to 2	
years, and a group of trees all	
die at the same time. Bark	
beetles commonly attack trees	
weakened by root disease	
infection.	
The incidence of root	
disease often relates to the	
amount of human activity in	
forest sites. Harvesting timber	
from infected stands often	
increases root disease activity.	
In addition, changes in forest	
structure brought about by	
logging have led to increases	
in tree species more prone to	
root disease.	
	Eigung 264 Boot
	Figure 264. Root disease killed
	saplings associated
	with stumps.
	Figure 265. Root
	disease infected
	trees are susceptible
	to bark beetle
	linfestations

Armillaria or Shoestring Root Rot Armillaria spp.

ponderosa pine,

Symptoms/

signs: The most diagnostic trait of *Armillaria* is the thick, fan-shaped mat of white mycelium in the cambium of roots and root crown. This may be accompanied



Figure 266. White mycelial fans are found under the bark of infested trees.

common in the Southwest. Other signs of the fungus include rhizomorphs, or black shoestring-like structures, on the outside of

infected roots. The fruiting bodies of this fungus are commonly produced in clusters but can be found singly. The caps are yellow brown to dark tan, with small scales on upper surface. The stem is stout, tapering upward to a ring of tissue just below the cap. The decay is light yellow, soft and spongy to stringy often containing numerous black zone lines.

Biology: Armillaria root decay spreads primarily by rhizomorphs and root contacts. Armillaria invades the bark and cambial region of roots and the root collar, killing roots and trees of all sizes. Wood decay follows cambial attack, and the wood serves as a source

Figure 267. Armillaria produces shoestring-like rhizomorphs.

of energy necessary for infection of new hosts. New infections are less common, occurring when mushrooms release windborne spores that germinate and colonize recently killed material. As with other root disease fungi, Armillaria can persist for decades in decaying wood in soil.

Armillaria often acts in conjunction with other secondary pests and pathogens. For example, *Dendroctonus*

may attack trees with this root disease.

Effects: Armillaria is the most common root disease in the Southwest and may account for up to 80 percent of the root disease conifer mortality in the region. There are over 10 species of Armillaria, some are virulent parasites while others are opportunists that act selectively on small or weak individuals such as those shaded by taller plants, defoliated by insects, attacked by other fungi, or weakened by drought.



Figure 268. Fruiting bodies of Armillaria spp.




Figure 269. Decay of Armillaria is soft and spongy.

Similar Insects and Diseases:

Other fungi such as *Fomitopsis pinicola* produce white mycelium beneath the bark of infected stumps, but the mycelium is not thick and fan-shaped like that produced by *Armillaria* spp.


References: 31, 90, 92, 112, 114

Annosus Root Rot Heterobasidion annosum (Fr.:Fr.) Bref. (Fomes annosus (Fr.:Fr.) Cooke)

Hosts

HOSIS:		
Symptoms/signs:	Fruiting bodies of annosu	s are grey-
brown on the upper surface	e with an undersurface that	t is chalky
white with pores. They are	e found on the underside of	f decayed roots
and slash, inside stumps (c	ommon in the Southwest),	or under the
duff at the base of infected	stumps and trees (althoug	h rare in the
Southwest). H. annosum le	eads to crown thinning and	d mortality or
windthrow, the latter of		
which can occur before		
aboveground symptoms		
are evident.		
Biology:		
Spores of <i>H. annosum</i>		
are produced on		
conks in decayed		
stumps or on roots of		
windthrown trees. The		
most common means		
		Figure 270.
		Fruiting bodies of H. annosum
		are often flat
		with the pores
		facing down.

Figure 271. Heterobasidion annosum is commonly found on underside of roots and inside stumps.



of initial entry of *H. annosum* into a site is via airborne spores that germinate on freshly cut stumps and basal wounds. Mycelium of H. annosum quickly colonizes the stump and grows into its roots. Transmission to adjacent trees occurs via root contacts. In live trees, the fungus decays woody root systems and then advances to the root collar where it may surface to the cambium and kill by girdling as in ponderosa pine, or progresses more slowly through roots to the stem

Although most conifers are susceptible to infection, there appear to be two types of annosus in North America, each of

pine but may also infect juniper. Like *Armillaria*, *H. annosum* is a common decayer of dead woody material as well as a pathogen.

Effects: This is one of the most important forest tree diseases in temperate zones worldwide. Damage to forests in North America has been most severe in the Southeast, where intensive

forestry has the longest history. In the Southwest, annosus root disease is the second most common root disease of conifers. It is found most often on true

areas where it is common in ponderosa pine. Infected trees are predisposed to attack by bark beetles.

Similar Insects and Diseases: The aboveground symptoms are similar to those caused by other root disease fungi.

References: 29, 53, 59, 73, 88, 92, 114

Figure 273. Wood discoloration is an early sign of annosus decay.

Tomentosus Root Disease Inonotus tomentosus (Fr.) Teng

Hosts:

Symptoms/signs: Fruiting bodies of *I. tomentosus* most commonly develop in August and September. They are small (usually less than 10 cm in diameter), mushroom shaped, and have a lower pore surface from which the spores are released. Although leathery, they are annual. The upper surface is yellow-brown to rust-brown and velvety, and becomes dark brown with age. The early stage of decay is characterized by a red-brown discoloration in the heartwood of roots. The late stage is also reddish brown but appears lighter because

of numerous small elongate pockets with pointed ends,

mycelium. This type of decay is known as a white pocket rot. The most distinctive characteristic of tomentosus rot is that a cross section



Figure 274. Leathery I. tomentosus fruiting body.

of an infected stem in advanced stages of decay has a honeycomb appearance.

Biology: Tomentosus root disease spreads primarily by root-to-root contact, however, infection by spores can occur through deep wounds in roots. There is no evidence that spores colonize stump surfaces as with annosus root disease. Diseased trees occur singly or in groups. The fungus spreads both outward in roots and up into the butt. Diseased roots have dead, decayed distal portions and red-brown, resin-soaked wood extending into the living portion and

the butt. Advanced decay develops slowly and expands to involve both deep-lying wood and sapwood near the cambium. Infected trees often die standing, but sometimes blow over while alive because of advanced decay in major roots or butt. *I. tomentosus* has been found to persist for more than 50 years in decaying wood in soil.

Effects: Although tomentosus root disease has been

there is little information available on the extent and damage from this disease. In other parts of the country, blowdown caused by tomentosus root disease has initiated spruce beetle (*Dendroctonus rufipennis*) outbreaks but that has not been reported here.

Similar Insects and Diseases: I. tomentosus and I. circinatus

Microscopically, *I. circinatus* can be differentiated from *I. tomentosus* based on hooked structures the latter does not produce. *I. tomentosus*

and *I. circinatus* are sometimes called false velvet-top fungi, because of the similarity of the conks of these fungi with that produced by *Phaeolus schweinitzii*, which is the real velvet-top fungus.



Figure 275. Red-brown discoloration of roots decayed by I. tomentosus.

References: 29, 92, 102, 110

Schweinitzii Root and Butt Rot Cow Patty, or Velvet Top Fungus Phaeolus schweinitzii (Fr.:Fr.) Pat. (Polyporus schweinitzii Fr.:Fr.).

Hosts:

pine, spruce

Symptoms/signs: Trees infected with *P. schweinitzii* do not typically show outward symptoms, so the presence of disease is usually not realized until a tree breaks or is windthrown due to loss in structural support. The fruiting bodies of *P. schweinitzii* often emerge through soil from decayed roots. They have a velvety

a yellow margin. It turns a dark red brown upon aging, is crumbly and can be mistaken for a cow pie at this point, hence the nickname

"the cow patty fungus." Incipient decay appears as a light yellow to pale reddish-brown color. As the decay advances, the color

the wood rapidly loses strength. In advanced stages of decay, the wood breaks up into large cubes that are crushed easily,

Figure 276. Fruiting structures of P. schweinitzii popping up from decayed roots.

and the butt portions of infected trees can be swollen.

Biology:

Spores likely initiate most *P. schweinitzii* infections. *P. schweinitzii* is unusual in that the fungus establishes itself in forest soil and enters trees through root tips. There is no root-to-root spread as used by other



Figure 277. Young specimen of P. schweinitzii.

1-2 m of the butt log, rarely extending much higher.

Effects: Schweinitzii root and butt rot is a major disease of older trees. It causes decay of the roots and lower stem and the resulting strength loss predisposes trees to windthrow and breakage.

Similar Insects and Diseases: The fruiting bodies of *I. tomentosus* and *I. circinatus* are similar to *Phaeolus schweinitzii* since they are all velvety on top when fresh and have a pore surface beneath. However, *P. schweinitzii* fruiting bodies are larger, greater than 15 cm in circumference, while the other two are about 5 cm maximum

References: 30, 92

Black Stain Root Disease Leptographium wageneri (Kendrick) MJ Wingfield

Hosts:

Symptoms/signs: Like other root diseases, black stain usually exhibits symptoms of gradual decline before tree death. In the early stages of decline, terminal growth is reduced and older needles become chlorotic. As the disease progresses, older needles are shed prematurely, new needles are somewhat stunted and yellow, and reduced internodal growth is evident on lateral branches. Very small trees may succumb quickly without exhibiting gradual decline symptoms.

Black stain root disease is distinguished from other root diseases by the dark-brown to purple-black discoloration in the sapwood of the lower bole and root collar. When observed in cross section, the black stain appears in arcs roughly concentric with the growth rings.




Figure 278. Black stain of piñon caused by Leptographium wageneri.

Biology: The Southwestern Region has two physiologically and morphologically distinct variants of black stain root disease. *Leptographium wageneri var. wageneri L. wageneri var. pseudotsugae* (Harrington et Cobb) causes black

but not in Arizona. A third variety *L. wageneri var. ponderosum* infects hard pines in California but has not been found in the Southwest.

Effects: Black stain root disease fungi grow in sapwood and plug tracheids, which prevents water transport and causes a wilting and rapid tree decline and death. Unlike *Armillaria* spp. and *H. annosum*, *L. wageneri* does not cause wood decay and dies with its host.

Black stain root disease affects groups of trees in distinct infection centers. Typical infection centers have trees in various stages of decline near the perimeter and dead trees in the interior where infection originated.

Similar Insects and Diseases: Blue stain fungi, which are often confused with black stain fungi, are usually a lighter color and typically are wedge-shaped in cross-section and they can discolor the entire sapwood radius.

References: 37, 92

Ganoderma Root Rot Artist's Conk Ganoderma applanatum (Pers.) Pat.

Hosts: Aspen

Symptoms/signs: Infected trees often break or fall before death, however, a fruiting body or conk can be found at the base of most infected trees. The tough semicircular conk is usually found near ground level. It has a brown upper surface with a white rim and an undersurface that is white and stains when touched or scratched. The latter trait allows for artistic expression that is permanent if the conk is allowed to dry. *G. applanatum* produces advanced whitemottled decay in the large roots and lower parts of the stem.

Biology: G. applanatum infection occurs at wounds and

the fungus attacks sapwood, heartwood, and cambium. The white-

Figure 279. Ganoderma applanatum conks are often found at the base of infected trees.




Figure 280. Trees often topple over due to severe decay of major roots.

mottled rot is usually concentrated in large roots (larger than 5 cm in diameter) and basal part of the stem, but can extend up into the trunk for a meter or more. The larger roots are believed to act as avenues of spread to new hosts.

Effects: *G. applanatum* is found in almost all aspen stands but is more abundant on moist sites with deep soils. Windthrow is common.

Similar Insects and Diseases: Although there are other root decay fungi of aspen, *G. applanatum* is the most prominent and the only one that produces a tough woody-conk at ground line in live trees in the Southwest.

References: 38, 92




Figure 281. Aspen stand with mortality caused by frost and drought.

Drought

when water loss by transpiration exceeds the rate of absorption from soil. Chronic injury results from long-term exposure to low water

supplies (e.g. in areas with poor soils) and is expressed by growth loss and increased susceptibility to parasitic fungi and insects. Acute injury occurs under

loss or death. Symptoms of drought damage include wilting, discoloration of foliage and premature leaf fall. Progression within the crown is from the top down and outside in, and



Figure 282. Western soapberry with branch dieback following dry spring.

the roots are usually the last portions to die. Symptoms can be similar to those of trees suffering from root disease, except that with root diseases, the roots die before the foliage. In the Southwest, drought symptoms have been observed in many trees including

soapberry.

Winter drought is particularly harmful in Southwestern upland forests and can render large areas vulnerable to bark beetles and

Figure 283. Aspen branch dieback during summer drought.

References: 82, 92

Frost

Frost damage of many tree species including aspen, locust, spruce, and oak has been observed throughout the Southwest. The tolerance of foliage and small branches to cold temperature depends on their "hardiness," which is closely correlated with the time of year and species. Temperatures just above freezing and short photoperiods induce frost hardiness. The sudden onset of below-freezing temperatures (frosts) in spring or early autumn can severely damage unhardened tissues, killing young shoots and leaves or needles.




Figure 284. Aspen leaves succumb to June frost.

As temperatures rise in late winter and early spring, plants deacclimate until, by the time growth begins they can no longer to

the time growth begins, they can no longer tolerate more than a few degrees of frost. Leaves and stems killed by spring frost are usually small and succulent at the time of injury. More mature leaves and stems can tolerate lower temperatures. After thawing, damaged



Figure 285. Newest and most sensitive growth of fir killed by frost.

reddish brown, dark brown, or black, depending on the species. Dead leaves and shoots break off or abscise during the next several weeks. New shoots and leaves begin to grow almost immediately and soon mask the early season damage.

References: 82,

92

Winter Injury




Figure 286. Needle bases of 1-year needles are green but tips are brown.

There are several different types of winter damage of conifers in the

develop in dormant evergreens during warm weather in late winter or early spring when needles transpire water from leaves and stems while the soil is cold or frozen. Roots

from cold soil and none

from frozen soil. Freeze damage occurs when needles no longer have enough moisture to survive normal nighttime temperatures. Freeze damage can also follow untimely deacclimation from temporary warm weather. Ponderosa pine and alligator juniper are the species most often observed with damage by these phenomena. Sometimes the result is trees that have needles with brown tips and green bases, and other times trees may have winter-killed needles that are completely brown regardless of age, but the branch and bud tissues are still green

and viable. In either case, trees recover during the growing season. Occasionally, severe freeze damage near the upper elevational limits of a particular tree species causes mortality or topkill.

References:

82, 92

Figure 287. Winter injury can be mistaken for mortality at a distance.

Red Belt

Red belt, another type of winter damage, is named because

bands varying from less than 20 to as much as 1,000 yards wide on slopes and benches. These events are associated with the sudden occurrence of warm, dry winds (e.g. Chinook winds) that produce a temperature inversion. A relatively thin layer of warm air arrives that cannot mix downward and continues to contact side slopes. Trees exposed to unseasonably warm air by day receive seasonably cold air at night. This alteration of warm and cold air exposure, along with the frozen condition of the soil, results in desiccation injury because daytime transpiration removes moisture from the needles more rapidly than roots in frozen soil can replace it. Later, older needles fall off, leaving only current year needles on living branches after bud-break.




Figure 288. A band of cold air gets trapped at a specific elevation around mountains.

Windthrow

Windthrow is most common when soils are shallow and/ or water saturated. Spring is very windy throughout much of the Southwest, and is the time when blowdown occurs in many root diseased infected areas. Blowdown is also prevalent in winter months when trees are loaded with snow.

More severe windthrow damage takes place with tornadoes and microbursts, which occasionally occur in northern Arizona. Tornadoes and microbursts both result in windthrown trees lying parallel to one another in the same direction as the storm. Tree species with shallow root systems (e.g. spruce) are more prone to windthrow.

Reference: 107

Figure 289.
There are typically decay or soil defects associated with windthrown trees.

Figure 290. Tornadoes and microbursts have been known to knock down a swath of trees.

Lightning

Lightning damage is thought to be responsible for approximately 2 percent of the annual tree mortality in the region. However, many more trees are slightly damaged than are killed by lightning. Killed trees are often blown apart at impact or soon colonized by bark beetles.

Damage is in the form of a long narrow furrow in which the bark and a thin layer of wood are blown out. Sometimes, but not always, lightning damage can result in internal decay that may provide cavity-nesting habitat for wildlife.

References: 114

Figure 291. Lightning struck trees are often attacked by bark beetles.

Figure 292. Many lightning struck trees survive for decades and are slowly decayed.

Salt Damage

The uptake of salts by the roots of roadside trees is a common problem wherever sodium chloride, calcium chloride, and magnesium chloride are applied to de-ice highways in the winter or for dust abatement in the summer. Most of the time symptoms are

remain green and trees rec	over. However, a surprising amount of
mortality may occur when	magnesium chloride or calcium chloride
	ent during or shortly before heavy
	rainstorms. Salts are leached off the road in heavy rains and travel into drainages where subsequent tree damage and mortality take place.
Figure 293. Salts applied for cause as much damage as the de-icing.	References: 6, 82, 92
Figure 294. Most tree species experience tip burn from salt damage.	

Burls

Burls are large woody, more or less hemispheric, swellings on the main stems of trees. Although some stem swellings are the result of pathogens, often the cause of burls is unknown. Burls may often bear many buds or sprouts.




Figure 295. The cause of burls is often unknown.

Ungulates




	Figure 298. I snapped thes saplings whi feeding on foliage after protective fer was removed	se le a nce
Figure 299. Ponderosa pine browsed by ungulates.		

Black Bear Ursus americanus

Black bears rip wide strips of bark from trees, often starting at about 1 m from ground line. They forage on the cambial zone in the lower bole by removing or peeling the bark with their claws and scraping the vascular tissue with their incisors. Trees are often girdled and die from one stripping, but sometimes survive. The wound left in partially girdled trees can lead to extensive decay. Bears also break branches on oak trees to access acorns.

In the White

Mountains of eastern Arizona, black bear strip the bark of southwestern white pine and subalpine

Figure 300. Black bears rip strips of bark from trees and feed on the cambium.

References: 46, 49, 93

Porcupine

Although porcupines have a variable diet during the growing

season, in the winter they chew through the outer bark of branches and boles (less than 2 centimeters in diameter) and feed on cambium. Coniferous trees are preferred and often girdled. Porcupine injury can be locally damaging because the animal tends to stay in a small area, killing the tops of many trees. Since porcupine populations tend to

form new leaders and grow to maturity.

Figure 301. Porcupine chew through the bark and feed on the cambium.

Sapsucker




Figure 302. Sapsuckers drill parallel rows of holes and feed on sap and insects caught in the sap.

Glossary

A

Abdomen: Posterior part of the insect's main body divisions.

Abiotic disorder: Interference of the normal functioning of a plant that is caused by some non-living factor (e.g., weather) and not by an organism (pathogen or insect); symptoms may be similar to a biotic disease.

Aecia: A cuplike structure bearing aeciospores in the rust fungi.

Adult: Full-grown, sexually mature insect; usually with wings in contrast to larvae which lack wings.

Alternate host: One of the two dissimilar plants infected by a heterocyclic fungus, i.e., rust fungi that use two hosts to complete a life cycle.

Arthropod: Largest phylum in the animal kingdom including insects, crabs, spiders, etc.

В

Basidiospore: Spore formed after sexual recombination has occurred.

Biotic: A living organism.

Blue stain: A bluish to grayish discoloration of sapwood of a killed tree caused by fungi that utilize the material without causing decay, has no effect on the strength of the wood.

Brown Rot: Brownish, dry, crumbly decay of wood caused by fungi

state; may occur in areas surrounded by sound wood. Decayed wood typically brakes into cubical shapes.

Boring-dust: Fragments of phloem or wood excavated by adult bark beetles as it bores through the outer bark; brown to tan, often found in small piles in bark crevices and around the base of a tree.

Brood: The offspring or next generation of young hatched from eggs laid by one series of parents, and that mature about the same time.



Callus: Healing tissue of trees on branches, boles, or roots that attempts to grow over scars or wounds.

Cambium: Layer of cells between xylem and bark, forms additional xylem and phloem elements.

Canker: Localized, dead, sometimes sunken, portion of the cambium and bark of branches or the bole of a tree, often caused by a pathogen. Many layers of dead callus folds surround some cankers as the tree repeatedly attempts to heal the dead portion, only to have the pathogen kill the callus folds.

Chlorosis: (adj. chlorotic) An unseasonable yellowing of the foliage, may occur in bands on needles.

Cocoon: A thin covering often largely of silk which an insect larva forms about itself and in which it passes the pupal stage.

Conk: Fruiting body of a wood-decay fungus, often hoof or bracket-shaped and perennial, although some are annual; other than a mushroom.

Context: Interior tissue of a fruiting body.

Cornicle: Dorsal tubular processes on the posterior portion of the abdomen; common on aphids.

Cortex: Rind or outer layer.

Crawler: The dispersal life stage of aphids and scale insects.

Crop tree: Tree left after thinning, often because of superior quality or species and lacking defect or disease.

D

Decay: Rot or destruction and decomposition of plant material by fungi; wood decay is typically separated into brown or white rot depending on the wood compounds consumed.

Declivity: A term used to describe the descending distal slope of beetle forewings. For example, Ips beetles have spines on their elytral declivity, while Dendroctonus species have a rounded elytral declivity.

Defoliator: An organism that causes damage to plant foliage.

Diapause: The dormant phase of some insects. Can be either facultative or obligatory.

Dioecious

plants, as in mistletoe.

Disease: A disturbance of a plant caused by a pathogen (not an insect) or environmental condition that interferes with the plant's normal growth, structure, or reproduction. See also abiotic disorder.

E

Egg niche: A small impression or incision created by female bark beetle in which an egg is deposited.

Elytra: Hardened forewings of beetles that serve to protect the functional posterior wings.

Endemic: Typical, nonepidemic population.

Endophytic: An organism living within tissues of a live plant, as in mistletoe.

Epidemic: Large-scale, temporary increase in an insect or disease population.

F

Fecundity: Ability to produce young.

Flagging: Dying or recently dead branches; contrasting in color with the normal green color of a living tree.

Frass: Solid insect excrement, usually in small pellets; many times mixed with boring dust.

Fruiting body: A general term for any fungal spore-producing structure. See also conk or mushroom.

Fungus: (plural=fungi) An organism incapable of producing its own food supply by photosynthesis, usually having microscopic thread-like feeding structures (hyphae) and reproductive spores. Fungi cause decay of plant material.

G

Gall: An abnormal growth of plant tissue, stimulated by insect or fungal activity.

Gallery: Engraving of cambium or bark caused by bark beetles.

Beetles create both egg galleries and feeding galleries.

Design of egg galleries are used to identify beetle species.

Genus: A group of closely related species. Similar genera are grouped in a family.

Girdle: The act of damaging the cambium completely around the circumference of the stem, root or branch, typically causing death of the tree or tissue beyond the point of girdling.

Gregariously: Insects tending to feed or remain in groups.

Grub

Н

Heartwood: Central mass of tissue in tree trunks, with no living cells and no longer functioning in water conduction, contributes to mechanical support.

Heart Rot: Decay, typically caused by fungi, that is

Heterocyclic: Fungi that requires two different hosts to complete their life cycle; i.e., some rust fungi.

Hibernacula: Shelters occupied during the winter by dormant insects, i.e., spruce budworm larvae overwinter in hibernacula.

Honeydew: Sugary liquid excretion of aphids or scale insects.

Host: A plant that is invaded by a parasite and from which the parasite obtains its nutrients.

Hypha: (plural=hyphae) a single branch of a mycelium.

Incipient decay: The early stage of decay in which degradation has not proceeded far enough to soften the wood or to cause a perceptible reduction in hardness. Such strength properties as toughness or impact strength are appreciably reduced in wood with incipient decay, but the damage is not normally visible to the naked eye.

Infection: The establishment of a parasite within a host plant.

Instar: The stage of an insect between successive molts.

Lanceolate: Spear-shaped, tapering at each end.

Larva: Immature form of an insect such as a caterpillar, grub, or maggot.

Lesion: A localized area of discolored, diseased tissue.

Latent infection: The state in which a host is infected with a pathogen but does not show any symptoms.

Life cycle: The stage or successive stages in the growth and development of an organism that occur between the appearance and reappearance of the same stage (e.g., spore) of the organism.

M

Metamorphosis: Change in form and structure between different life stages of insects. Some insects have incomplete metamorphosis (egg, nymph, adult), while others have complete metamorphosis (egg, larva, pupa, adult).

Midge

Molt: Process of shedding the exoskeleton, the insect "skin."

Monogamous: Mating with only one individual.

Mushroom: Fleshy fruiting body of a fungus that has either gills or pores.

Mycangia: Structure found on some insects used for the transportation of symbiotic fungi from one host to another.

Mycelium: (hyphae) The vegetative feeding structure of a fungus.

Mycelial fan: a fan-shaped mycelial mat forming under the bark and wood of roots or lower trunks; associated with *Armillaria* spp.

Mychorrhizae: Association between plant rootlets and specialized nutrient and water uptake.

N

Necrotic: Dead plant cells or tissues.

Nuptial chamber: Mating site.

Nymph: Immature form of an insect resembling the adult, except for

incomplete wing development.

0

Obligate parasite: An organism capable of existing only as a parasite on a live host.

Overwinter: Term used to describe what life stage an insect passes the winter in.

Oviposit: Lay or deposit eggs.

Ovipositor: Egg-laying apparatus.

P

Parasite: An organism living in or on another living organism (host) and obtaining its food from the latter. May be harmful (i.e.,

Parthenogenetic: Reproducing by eggs that develop without being fertilized.

Pathogen: An organism or virus capable of causing disease on a host. Many pathogens of forest trees are fungi.

Perennial: Lasting for several years or more.

Pheromone: Chemicals produced by one individual to affect or alter the behavior of another individual of the same species.

Many bark beetle species produce both aggregation and anti-aggregation pheromones, which mediate the colonizing behavior of other beetles.

Phloem: Vascular tissue that conducts synthesized foods through the plant, located adjacent to the outside of the cambium in trees, essentially the inner bark.

Pitch-tubes: A tubular mass of resin mixed with bark, wood borings, and insect excrement that form on the surface of the bark at beetle entrance holes.

Plumed: Feather-like.

Polygamous: Mating with several individuals.

Pronotum: The dorsal plate of the prothorax.

Proleg

insect larvae but not in the adult.

Prothorax: The anterior of the three thoracic segments.

Punk knot

beginning of a new conk, also called blind conk, reliable indicator of the presence of some wood decay fungi, i.e., *Phellinus pini*.

Pupa: Inactive stage of an insect, a transitional stage from larva to adult.

R

Radial growth: One-half the diameter growth of a tree.

Resinosis

Rhizomorph: A specialized form of mycelium produced by certain fungi (e.g. *Armillaria* spp.) consisting of several strands of hyphae twisted together so as to be rootlike, and covered with a protective sheath.

Rot: Decay or deterioration of organic material through the enzymatic activity of microorganisms.

Rust: A disease caused by one of the rust fungi; highly specialized pathogens that are obligate parasites, which frequently require two different types of hosts to complete their life cycles (heterocyclic), and produce orange, yellow, or brown spores.



Saprophyte: An organism living in and getting its food from dead organic material.

Sapwood: Outer region of xylem of tree trunks, containing some living cells and functioning in water conduction, food storage and mechanical support.

Scales: A small thin, plate-like structure found on the wings of Lepidoptera.

Serrate: Saw-toothed edge.

Shootborer: Insects that feed in the pith of shoots.

Sign: The pathogen or its parts or products seen on a host plant.

Slash: Debris such as logs, bark, and branches left after cutting timber.

Spore: The reproductive unit of fungi consisting of one or more cells, analogous to the seed of green plants.

Sporophore: Fruiting body of a fungus that produces spores (i.e., mushroom or conk).

Stage: A distinguishable period of growth and development of an insect, i.e., the larval stage of an insect.

Stomata: Breathing pore in the epidermis (outer layer) of a plant.

Subcortical: Below the bark.

Sunscald: Death of bark due to sudden exposure to sun. This is common in mature aspen where neighboring trees have been removed.

Symbiotic: An intimate association between two species, which

Symptom: The external and internal reactions or alterations of a plant as a result of a disease or insect.

Systemic: Throughout the plant host tissues, usually surviving from year to year.

Т

Target canker: Perennial canker produced by a fungus which invades live tissue during the host dormant season, and survives in dead material during the host growing season.

Terminal growth: Height growth of a tree.

Terpenes: Unsaturated hydrocarbons occurring in plant oils and resins, common in conifers.

Thorax: The body region of an insect behind the head, which bears the legs and wings.



Vector: An organism such as an insect that transmits a pathogen.



White rot: White to tan decayed wood, caused by fungi decomposing both lignin and cellulose; may occur in areas

or spongy in texture, and may contain distinctive pockets. In advanced stage, may create a large hollow in a tree.

- **Windthrown**: Trees uprooted by wind. The term is also sometimes applied to trees whose stems are snapped by the wind.
- **Witches' brooms**: Abnormal proliferations of shoots or branches caused by a pathogen (e.g., from dwarf mistletoe).
- **Woodborer**: Diverse group of insects that feed at least in part in the wood. Many of these insects begin their larval stage in the cambium before constructing galleries in the xylem (wood).



Xylem: Vascular tissue that conducts water and mineral salts, taken in by roots, throughout the plant, essentially the woody part of the stem or trunk.



Zone line: Narrow black to brown lines in decaying wood formed by fungi.

Photography and Illustration Credits

- Robert L. Anderson (USDA Forest Service, www.forestryimages. org); 188
- Ronald F. Billings (Texas Forest Service, www.forestryimages.org); 45, 149
- Bob Cain (USDA Forest Service, Rocky Mountain Region); 190
- Clemson University (USDA Cooperative Extension Slide Services, www.forestryimages.org); 44
- Conklin, David A.; 2, 15, 37, 191, 194-195, 198, 202, 204, 218-224, 230-235, 238, 240-252, 254-257, 261, 263, 295
- Whitney Cranshaw (Colorado State University, www.forestryimages. org); 61-62
- Tom DeGomez (University of Arizona, Cooperative Extension); Cover-1,133
- Jerald E. Dewey (USDA Forest Service, www.forestryimages.org); 13, 24, 26, 46, 80
- Brian Geils (USDA Forest Service, Rocky Mountain Research Station); 228-229
- John H. Ghent (USDA Forest Service, www.forestryimages.org); 274
- Kennenth E. Gibson (USDA Forest Service, www.forestryimages. org); 83c, 104, 122
- Thomas E. Hinds (USDA Forest Service, Rocky Mountain Research Station); 176, 210-211, 216, 225,
- Edward Holsten (USDA Forest Service, www.forestryimages.org); 50, 57, 170
- Gerald J. Lenhard (Louisiana State University, www.forestryimages. org); 43
- R. Ladd Livingston (Idaho Department of Lands); 83a, 117, 150
- Ann Lynch (USDA Forest Service, Rocky Mountain Research Station); 41
- David McComb (USDA Forest Service, www.forestryimages.org); 5

- Roy Morales (USDA Forest Service, Carson NF); 49
- Minnesota DNR (Minnesota Department of Natural Resources Archives, www.forestryimages.org); 51
- A. Steven Munson (USDA Forest Service, www.forestryimages.org); 118
- Donald Owen (California Department of Forestry and Fire Protection, www.forestryimages.org); 130
- Jeff Schalau (University of Arizona, Cooperative Extension); 92
- John Schwant (USDA Forest Service, www.forestryimages.org); 189
 - 174, 178, 183-184, 275, 284, 294
- Scott Tunnock (USDA Forest Service, www.forestryimages.org); 76-77, 144
- Michael R. Wagner (Northern Arizona University); Cover-3, 20-21, 23, 25, 27-28, 30-32, 48, 75, 78, 151-157, 161, 164
- USDA Forest Service, Intermountain Region Archives (www. forestryimages.org); 3, 10, 162
- USDA Forest Service, Northern Region; 83b (R1-03-08)
- USDA Forest Service, Rocky Mountain Region Archives (www. forestryimages.org); 4, 7, 8, 11, 29, 52, 105, 107, 127
- USDA Forest Service, Southwestern Region; Cover, 1, 3-4, 6, 9, 14, 16-19, 33-36, 38-40, 42, 47, 53-56, 58-60, 63-71, 73-74, 79, 81-82, 84-91, 93-101, 103, 106, 108-116b, 119-121, 123-126, 128-129, 131-132, 134-143, 145-148, 158-160, 163, 165-169, 171-173, 175, 177, 179-182, 185-187, 192-193, 196-197, 199-201, 203, 205-209, 212-215, 217, 226-227, 236-237, 253, 258-260, 262, 264-273, 276-283, 285-293, 296-302

Station: 239

Instructions for Submitting Insect and Disease Specimens

shown on next page). The following procedures for collecting and shipping specimens should be used:

Collecting

- 1. Adequate material should be collected
- 2. Adequate information should be recorded, including:
 - a. location of collection
 - b. when collected
 - c. who collected the specimen
 - d. host description (species, age, condition, etc.)
 - e. area description (forest type, site conditions, etc.)
 - f. unusual conditions (frost, poor drainage, etc.)
- 3. Personal opinion of the cause of the problem may be helpful.

Packing

- 1. Larvae and other soft-bodied insects should be shipped in small screw-top vials or bottles containing at least 70 percent isopropyl (rubbing) alcohol. Make sure bottles are well sealed.
- 2. Pupae and hard-bodied insects may be shipped either in alcohol or in small boxes. Specimens should be placed between layers of tissue paper in the boxes. Pack carefully and make sure there is little movement of material within the box. Do not pack insects in cotton.

3. Needle or foliage diseases: Do not ship in plastic bags as condensation can become a problem. Use a paper bag or wrap in newspaper. Pack carefully and make sure there is little movement within the box.

Mushrooms and conks: Do not ship in plastic bags.

Either pack and ship immediately or air-dry and pack.

To pack, wrap specimens in newspaper and pack into a shipping box with more newspaper. If on wood, include some of the decayed wood.

Federal Forest Health Zone Offices

Arizona Zone Forest Health 2500 South Pine Knoll Drive Flagstaff, AZ 86001

New Mexico Zone Forest Health 333 Broadway Blvd., SE Albuquerque, NM 87102

State Insect and Disease Specialists

Arizona

Forest Health Program P.O. Box 15018 Northern Arizona University Flagstaff, AZ 86011

New Mexico

NMSU Coop. Ext. Forest Entomologist New Mexico EMNRD, Forestry Division P.O. Box 1948 Santa Fe, NM 87504

Selected References

- 1. Amman, G. D., M. D. McGregor, and R. E. Dolph, Jr. 1985. Mountain pine beetle. USDA Forest Service, Forest Insect
- 2. Barr, B. A., D. L. Hanson, and C. S. Koehler. 1978. Red turpentine beetle a pest of pines. University of California,
- 3. Beatty, J. 1979. Forest insect and disease conditions in the Southwest 1979. USDA Forest Service, Southwestern Region Publication 80-7, 29 p.
- 4. Beckwith, Roy C. 1973. The large aspen tortrix. USDA
- Blake, E. A., M. R. Wagner, and T. W. Koerber. 1985.
 Insects destructive to the ponderosa pine cone crops in northern Arizona. In: Proc. Conifer Tree Seed in the Inland Mountain West Symposium, Aug. 5-6, Missoula, MT. USDA Forest Service, Intermountain Research Station: 238-242.
- 6. Bolander, P., ed. 1999. Dust palliative selection and application guide. Project Report. 9977-1207-SDTDC. San Dimas, CA: USDA Forest Service, San Dimas Technology and Development Center. 20 p.
- 7. Borden, J. H., and M. McClaren. 1970. Biology of Cryptoporus volvatus (Peck) Shear (Agaricales, Polyporaceae) in southwestern British Columbia: distribution, host species, and relationship with subcortical insects. Syesis, Vol. 3: 145 154.
- 8. Cain, R., J. Cota, and C. Ward. 1990. Conifer pests in New Mexico. Rev. ed. (Albuquerque, NM): USDA Forest Service, Southwestern Region. 48 p.
- 9. Cain, R., and D. Parker. 1998. Conifer pests in New Mexico. Rev. ed. (Albuquerque, NM): USDA Forest Service, Southwestern Region. 50 p.

- 10. Carolin, V. M., and J. A. E. Knopf. 1968. The pandora
- 11. Childs, T. W., K. R. Shea, and K. L. Stewart. 1971. Elytroderma disease of ponderosa pine. USDA Forest
- 12. Conklin, D. A. 2000. Dwarf mistletoe management and forest health in the Southwest. USDA Forest Service, Southwestern Region, unnumbered report. 30 p.
- Corn, M. L., E. H. Buck, J. Rawson, and E. Fischer. 1999.
 Harmful nonnative species: issues for Congress. National Council for Science and the Environment. Report RL30123. 50 p.
- Coulson R. N. and J. A. Witter. 1984. Forest Entomology: Ecology and Management. John Wiley & Sons, New York, NY. 669 p.
- Cranshaw, W., D. Leatherman, W. Jacobi, B. Kondraitieff, and L. Mannix. 2000. Insects and Diseases of Woody Plants of the Central Rockies. Bulletin 506A. Colorado State University Cooperative Extension. Fort Collins, 283 p.
- 16. Dahms, C. W. and B. W. Geils, tech. eds. 1997. An assessment of forest ecosystem health in the Southwest. General Technical Report RM-GTR-295. Fort Collins, CO: USDA, Forest Service, Rocky Mountain Forest and Range Experiment Station. 97 p.
- 17. DeMars, Jr., C. J., and B. H. Roettgering. 1982. Western pine beetle. USDA Forest Service, Forest Insect and Disease
- Dunbar, C. S. and M. R. Wagner. 1990. Distribution
 (Hymenoptera: Diprionidae) in the United States and Canada. Entomol. News 101: 266-272.

- 19. Fellin, D. G. and J. E. Dewey. 1982. Western spruce budworm. USDA Forest Service, Forest Insect and Disease
- 20. Ferrell, G. T. 1986. Fir engraver. USDA Forest Service,
- Filip, G. M., and Schmitt, C. L. 1990. RX for Abies:
 Washington. USDA Forest Service, GTR-PNW-252. 34 p.
- 22. Funk, A. 1985. Foliar Fungi of Western Trees. Canadian
- 23. Furniss R. L. and V. M. Carolin. 1977. Western Forest Insects. USDA Forest Service, Miscellaneous Publication No. 1339. 654 p.
- 24. Furniss R. L. and J. B. Johnson. 2002. Field Guide to the Bark Beetles of Idaho and Adjacent Regions. University of Idaho. 125 p.
- Geils, B. W., J. Cibrian Tovar, and B. Moody, tech. coords.
 2002. Mistletoes of North American conifers. Gen. Tech.
 Rep. RMRS-GTR-98. Ogden, UT: USDA, Forest Service,
 Rocky Mountain Research Station. 123 p.
- Geils, B. W.; Conklin, D. A.; Van Arsdel, E. P. 1999. A preliminary hazard model of white pine blister rust for the Sacramento Ranger District, Lincoln National Forest. USDA Forest Service, RMRS-RN-6. 6 p.
- 27. Gilbertson, R. L. 1974. Fungi That Decay Ponderosa Pine. Tucson: University of Arizona Press. 197 p.
- 28. Gilbertson, R. L., and M. Rosemeyer. 1986. Comandra blister rust on Mondell pine in Arizona. Arizona Plant Pathology, Unnumbered Bulletin. Tucson, AZ; Dept. of Plant Pathology, University of Arizona. 3 p.

- 29. Gilbertson, R. L., and L. Ryvarden. 1986. North American
- 30. Gilbertson, R. L., and L. Ryvarden. 1987. North American
- 31. Hagle, S. K.; Byler, J. W. 1994. Root diseases and natural disease regimes in forest of western U.S.A. In: Johansson, M. and J. Stenlid, eds. Proceedings of the Eighth International Conference on Root and Butt Rots; 1993

 August 9-16; Wik, Sweden and Haikko, Finland. Uppsala, Sweden: Swedish University of Agricultural Sciences: 606-617.
- 32. Hagle, S. K., S. Tunnock, K. E. Gibson, and C. J. Gilligan. 1987. Field Guide to Diseases and Insect Pests of Idaho and Montana Forests. USDA Forest Service, Northern Region. 123 p.
- 33. Hawksworth, F. G. 1961. Dwarf mistletoe of ponderosa pine in the Southwest. Tech. Bull. 1246. USDA, Forest Service, Rocky Mountain Forest and Range Experiment Station. 13 p.
- 34. Hawksworth, F. G., and B. W. Geils. 1990. How long do mistletoe-infected ponderosa pines live? Western Journal of Applied Forestry. 5(2): 47-48.
- 35. Hawksworth, F. G. and R. F. Scharpf. 1981. Phoradendron7p.
- Hedlin, A. F., H. O. Yates, D. C. Tovar, B. H. Ebel, T.W. Koerber, and E. P. Merkel. 1980. Cone and Seed Insects of North American Conifers. Canadian Forestry Service. 122 p.
- 37. Hessburg, P. F., D. J. Goheen, and R. V. Bega. 1995. Black stain root disease of conifers. Forest Insect and Disease

- 38. Hinds, T. E. 1985. Diseases. In: DeByle, N. V., Winokur, R. P., eds. Aspen: Ecology and Management in Western United States. Tech. Rep. RM-119. Fort Collins, CO: USDA Forest Service, Rocky Mountain Forest and Range Experiment Station: 87-106.
- 39. Hinds, T. E. and M. G. Ryan. 1985. Expansion of sootybark and ceratocystis cankers on aspen. Plant Disease 69:842-844.
- Holsten, E. H., P. E. Hennon, L. Trummer, and M. Schultz.
 2001. Insects and Diseases of Alaskan Forests. USDA
 Forest Service, Alaska Region, R10-TP-87. 242 p.
- 41. Holsten, E. H., R. W. Their, A. S. Munson, and K. E. Gibson. 1999. The spruce beetle. USDA Forest Service, Forest
- 42. Itami, J. K. and T. P. Craig. 1989. Life history of Styloxus bicolor (Coleoptera: Cerambycidae) on Juniperus monosperma in northern Arizona. Annals of the Entomological Society of America 82: 582-587.
- 43. Jennings, D. T. and R. E. Stevens. 1982. Southwestern pine tip moth. USDA Forest Service, Forest Insect and Disease
- 44. Johnson, D. W., J. S. Beatty, and T. E. Hinds. 1995. Cankers on western quaking aspen. USDA Forest Service, Forest
- 45. Johnson W. T. and H. H. Lyon, 1988. Insects that feed on trees and shrubs, 2nd ed. Cornell University Press, Ithaca, NY. 556 p.
- Kanaskie, A., J. Chetock, G. Irwin, D. Overhulser. 1990.
 Black bear damage to forest trees in northwest Oregon 1988-1989, Pest Management Report 90-1, Oregon Department of Forestry, Salem, OR.

- 47. Keen, F. P. 1958. Cone and seed insects of western forest trees. USDA Forest Service, California Forest and Range Exp. Stn. Tech. Bull. 1169. 169 p.
- 48. Kegley, S. J., R. L. Livingston, and K. E. Gibson. 1997. Pine engraver, Ips pini (Say), in the western United States.

 7 p.
 - and S. Ferguson. 1998. Impacts of live canopy pruning on implications for black bear tree selection. Forest Ecology and Management. 109(1-3):51-56.
- Linnane, J. P., comp. 1984. Annual Southwestern Region pest conditions report-1983. USDA Forest Service, Southwestern Region. Albuquerque, NM. R3 84-13. 13 p.
- Lightle, P. C. and S. R. Andrews. 1968. Red rot in residual ponderosa pine stands on the Navajo Indian Reservation.
 USDA Forest Service, Rocky Mountain Research Station.
 Res. Pap. RM-37. 12 p.
- Livingston, R. L. 1979. The pine engraver beetle in Idaho, life history, habits and management recommendations.
 Idaho Department of Lands, Forest Insect and Disease Control Report 79-3. 7 p.
- 53. Livingston, W. H., A. C. Mangini, H. G. Kinzer, and M. E. Mielke. 1983. Association of root diseases and bark beetles (Coleoptera: Scolytidae) with Pinus ponderosa in New Mexico. Plant Disease 67: 674-676.
- 54. Lynch, A. M. 2003. Spruce aphid in high elevation habitats in the Southwest. p. 60-63. In: S. L. C. Gosbroke, K.W. Gottschalk, editors, Proceedings of the USDA interagency research forum on gypsy moth and other invasive species, 2002. USDA Forest Service, GTR-NE-300.

- 55. Lynch, A. M. 2004. Fate and characteristics of Picea damaged by Elatobium abietinum (Walker) (Homoptera: Aphididae) in the White Mountains of Arizona. Western North American Naturalist. 7-17.
- Lynch, A. M. and Fitzgibbon, R. A. 2005. Biology and USA, submitted for publication.
- 57. Lynch, A. M., J. L. Wilson, and M. S. Frank. In press. Elatobium abietinum (Walker) populations reach epidemic levels on Picea engelmannii Parry in the interior southwestern United States. Journal of Agricultural and Forest Entomology.
- 58. Maffei, H., and J. Beatty. 1988. Changes in the incidence of dwarf mistletoe over 30 years in the Southwest. Van der Kamp, B. J. ed., Proceedings 36th Western International Forest Disease Work Conference; 1988 September 19-23; Park City, UT. Vancouver, BC: University of British Columbia: 88-90.
- 59. Maijala, P., T. C. Harrington, and M. Raudaskoski. 2003. A peroxidase gene family and gene tree in Heterobasidion and related genera. Mycologia 95:209-221.
- 60. Massey, C. L., Lucht, D. D., and J. M. Schmid. 1977. Roundheaded pine beetle. USDA Forest Service, Forest

116. 6 p.

62. McCambridge, W. F. 1974. Pinyon needle scale. USDA

Neodiprion edulicolis Ross. USDA Forest Service, Forest

- 64. McMillin, J. D., K. E. Allen, D. F. Long, J. L. Harris and J. F. Negrón. 2003. Effects of western balsam bark beetle Journal of Applied Forestry 18: 259-266.
- McMillin, J. D., G. Hengxiao, M. R. Wagner & X. Long. 1996. Host density and elevation effects on the spatial southwestern China. Forest Ecology and Management 86: 151–161.
- 66. McMillin, J. D., and M. R. Wagner. 1989. Assessing the impacts of foliage-feeding insects on timber and scenic beauty of ponderosa pine: a methodological approach. p. 51–59. In: (A. Tecle, W. W. Covington & R. H. Hamre, tech. coor.) Multiresource Management of Ponderosa Pine Forests. USDA Forest Service GTR RM–185.
 - dynamics. p. 333–361. In: (M. R. Wagner & K. F. Raffa, Academic Press, Orlando, FL.
- 68. McMillin, J. D. and M. R. Wagner. 1997. Chronic performance and host plant quality. Oikos 79: 357–362. host plant versus natural enemies on the spatial distribution Diprionidae). Ecological Entomology 23: 397–408.
- 70. Mielke, J. L. 1943. White pine blister rust in western North America. Yale Univ. School of Forestry. Bulletin No. 52. 155 p.
- 71. Millar, C. S. 1984. Lophodermella species on pines. p. 45-55. In: Recent Research on Conifer Needle Diseases.

- Ed. Glenn W. Peterson. USDA Forest Service, General Technical Report GTR-WO 50.
- 72. Niwa, C G., R. E. Sandquist and 19 others. 2001.
 Invertebrates of the Columbia River basin assessment area.
 (T. M. Quigley, Ed.) Interior Columbia Basin Ecosystem
 - Service, General Technical Report PNW-GTR-512, Portland, OR. 74 p.
- 73. Otrosina, W. J.; Scharpf, R. F., technical coordinators. 1989. Proceedings of the Symposium on Research and Management of Annosus Root Disease (Heterobasidion annosum) in Western North America; April 18-21, 1989; Monterey, CA. Gen. Tech. Rep. PSW-116. Berkeley, CA:
 - USDA Forest Service. 177 p.
- 74. Parker, D. L. 1980. Mountain pine beetle, Dendroctonus ponderosae Hopkins, in ponderosa pine, Kaibab Plateau, Arizona. USDA Forest Service, Southwestern Region, Integrated Pest Management Guide 80-8. 12 p.
- 75. Parker, D. L. 1991. Integrated pest management guide: engraver, Ips pini (Say), in ponderosa pine. USDA Forest Service, Southwestern Region, 91-8. 17 p.
- 76. Parmeter, J. R., Jr. 1978. Forest stand dynamics and ecological factors in relation to dwarf mistletoe spread, impact, and control. p. 16-30. In: Proceedings of the Symposium on Dwarf Mistletoe Control Through Forest Management. Robert F. Scharpf and John R. Parmeter, Jr., Tech. Coords., April 11-13, 1978, Berkeley, CA: USDA Forest Service, General Technical Report PSW-31. 190 p.
- 77. Peterson, Glenn W. 1982. Dothistroma needle blight of

- 78. Peterson, R. S. 1967. Studies of juniper rusts in the West.
- 79. Peterson, R. S. 1968. Limb rust of pine: the causal fungi. Phytopathology 58: 309-315.
- 80. Rindge, F. H. 1967. A new species of Nepytia from the southern Rocky Mountains. New York Entomological Society LXXV: 74-76.
- 81. Schalau, J. 2003. Cypress bark beetles. The University of Arizona College of Agriculture and Life Sciences Publication AZ1316. 2 p.
 - Coast Conifers. Agricultural Handbook 521. Washington, DC, USDA Forest Service. 199 p.
- 83. Scharpf, R. F., and R. V. Bega. 1981. Elytroderma disease reduces growth and vigor, increases mortality of jeffery pines at Lake Tahoe Basin, California. USDA Forest Service. PSW-155. 6 p.
- 84. Scharpf, R. F., and Hawksworth, F. G. 1974. Mistletoes on hardwoods in the United States. USDA Forest Service,
- 85. Schmid, J. M., and D. D. Bennett. 1988. The North Kaibab Pandora Moth Outbreak, 1978-1984. USDA Forest Service, GTR-RM-153, 19 p.
- Schmid, J. M. and R. H. Frye. 1977. Spruce beetle in the Rockies. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, General Technical Report RM-49. 38 p.
- 87. Schmid, J. M, J. C. Mitchell, K. D. Carlin, and M. R. Wagner. 1984. Insect damage, cone dimensions, and seed production in crown levels of ponderosa pine. Great Basin Naturalist: 575-578.

88. Schmitt, C. L., J. R. Parmeter, and J. T. Kliejunas. 2000. Annosus root disease of western conifers. USDA Forest

p.

- 90. Shaw, C. G., III and G. A. Kile. 1991. Armillaria Root Disease. USDA Forest Service, Agriculture Handbook No. 691. 233 p.
- 91. Shepperd, Wayne D. and Fairweather, M. L. 1994. Impact of large ungulates in restoration of aspen communities in a southwestern ponderosa pine ecosystem. In: Covington, W. Wallace; DeBano, Leonard F.; tech. coords. Sustainable Ecological Systems: Implementing an Ecological Approach to Land Management. 1993 July 12-15; Flagstaff, Arizona. GTR-RM-247. Fort Collins, CO: USDA, Forest Service, Rocky Mountain Forest and Range Experiment Station.
- 92. Sinclair, W. A., H. H. Lyon, and W. T. Johnson. 1987.
 Diseases of Trees and Shrubs. Cornell University Press. 574
 p.
 - Characteristics Around Middens of Mount Graham Red Squirrels. Thesis submitted to the faculty of the School of Renewable Natural Resources, University of Arizona. 78 p.
- 94. Smith, R. H. Red turpentine beetle. USDA Forest Service,
- 95. Sower, L. L., G. E. Daterman, and C. Sartwell. 1989.
 Biology and impact of western pine shoot borer, Eucosma sonomana, in the western United States. p. 102-109. In:
 Proceedings of the IUFRO working group on Insects
 Affecting Reforestation, XVIII International Congress of Entomology, 1988. Vancouver, British Columbia, Canada.

- Stevens, R. E., J. W. Brewer, and D. T. Jennings. 1978. Life history and habits of Coleotechnites edulicola (Gelechiidas)
 Lepidopterists Society. 32(2): 123-129.
- 97. Stevens, R. E. and D. T. Jennings. 1977. Western pine shoot borer: A threat to intensive management of ponderosa pine in the Rocky Mountain area and the Southwest. USDA Forest Service, GTR-RM 45. 8 p.
- 98. Swetnam, T. W., and A. M. Lynch. 1989. A tree-ring reconstruction of western spruce budworm in the southern Rocky Mountains. Forest Science. 35(4):962-986.
- 99. Swetnam, T. W. and A. M. Lynch. 1993. Multicentury, regional-scale patterns of western spruce budworm outbreaks. Ecological Monographs. 63(4):399-424.
- 100. Tainter, F. H. and F. A. Baker. 1996. Principles of Forest Pathology. John Wiley & Sons, Inc. 805 p.
- 101. Tinnin, R. O. 1984. The effect of dwarf mistletoe on forest community ecology. p. 117-122. In: Frank G. Hawksworth and Robert F. Scharpf, Tech. Coords., Biology of Dwarf Mistletoes: Proceeding of the Symposium, 1984. USDA Forest Service, GTR-RM-111.
- 102. Tkacz, B. M. 1991. Survival of Inonotus tomentosus in spruce stumps after logging. Plant Disease 75(8): 788-790.
- 103. USDA Forest Service. 1985. Insects and diseases of trees in the South. USDA Forest Service, GTR-R8-5. 98 p.
- 104. USDA Forest Service. 1986. Tiger moth. USDA Forest Service, Southwestern Region, Fact Sheet. 2 p.
- 105. USDA Forest Service. 2002. Forest insect and disease conditions in the Southwestern Region, 2001. USDA Forest Service, Southwestern Region. Albuquerque, NM. R3-02-01. 17 p.

- 106. Van Arsdel, E. P., and B. W. Geils. 2004. The Ribes of Colorado and New Mexico and their rust fungi. FHTET-04-13. Fort Collins, CO: USDA Forest Service, Forest Health Technology Enterprise Team. 32 p.
- 107. Vasiloff, S. 1997. Microburst prediction and detection.N.O.A.A. Western Region Technical Attachment. No. 97-21.2 p.
- Wagner, M. R. 2001. Insects and diseases of southwestern forests. Lecture notes compiled by Michael R. Wagner, Northern Arizona University. 159 p.

5 p.

- 110. Whitney, R. D. 1977. Polyporus tomentosus root rot of conifers. Forestry Technical Report No. 18. Canadian Forestry Service; 12 p.
- 111. Wickman, B. E., R. R. Mason, and G. C. Trostle. 1981.
- Williams, R. E., C. G. Shaw, III, P. M. Wargo, and W.
 H. Sites. 1989. Armillaria Root Disease. USDA Forest
- 113. Wilson, J. L. and A. M. Lynch. 1998. Spruce aphid in the Southwest. USDA Forest Service, Southwestern Region.
- 114. Wood, R. E. 1983. Mortality caused by root diseases and associated pests on six national forests in Arizona and New Mexico. USDA Forest Service, Southwestern Region, Albuquerque, New Mexico. R-3 83-13. 31 p.
- 115. Wood, S. L. 1982. The bark and ambrosia beetles of North and Central America (Coleoptera: Scolytidae), a taxonomic monograph. Great Basin Naturalist Memoirs. Number 6. 1359 p.

- 116. Yohem, K. H., G. B. Cummins, and R. L. Gilbertson.1985. Revised list and host index of Arizona rust fungi.Mycotaxon. 23(2): 451-468.
- 117. Zabel, R. A. and J. J. Morrell. 1992. Wood Microbiology: Decay and Its Prevention. Academic Press, Inc. 476 p.

Host Index

Abies concolor
Branches & Terminals
True Fir Dwarf Mistletoe Arceuthobium abietinum 177
White Fir True Mistletoe Phoradendron pauciflorum 184
Foliage
Douglas-Fir Tussock Moth <i>Orgyia pseudotsugae</i>
Western Spruce Budworm Choristoneura occidentalis 8
White Fir Needleminer <i>Epinotia meritana</i> 16
Roots
Annosus Root Rot Heterobasidion annosum 193
Stem
Bark Beetles & Wood Borers Ambrosia Beetles
Decays
Indian Paint Fungus <i>Echinodontium tinctorium</i> 138 Red Ring Rot <i>Phellinus pini</i>
Carpenter Ants <i>Camponotus</i> spp
Termites
Rusts
Fir Broom Rust
Melampsorella caryophyllacearum
Abies lasiocarpa
Branches & Terminals
Douglas-Fir Dwarf Mistletoe Arceuthobium douglasii 177
Foliage
Nepytia janetae

Stem	
Bark Beetles & Wood Borers	
Ambrosia Beetles9	2
Western Balsam Bark Beetle Dryocoetes confusus7	9
Decays	
Indian Paint Fungus Echinodontium tinctorium 13	8
Rusts	
Fir Broom Rust Melampsorella	
caryophyllacearum	1
Acacia spp.—acacia	
Branches & Terminals	
Desert Mistletoe <i>Phoradendron californicum</i> 18	34
Acer spp.—maple	
Stem	
Bark Beetles & Wood Borers	
Ambrosia Beetles9	12
Alnus tenuifolia—thinleaf alder	
Foliage	
Alder Flea Beetle Altica ambiens	5
Fall Webworm <i>Hyphantria cunea</i> 2	29
Stem	
Cankers	
Cytospora Canker Cytospora chrysosperma15	52
Aphrophora spp	3
Arbutus arizonica—Arizona madrone	
Foliage	
Fall Webworm <i>Hyphantria cunea</i>	29
Cercidium spp.—palo verde	
Branches & Terminals	
Desert Mistletoe <i>Phoradendron californicum</i> 18	34

Cercocarpus spp.—mountain mahogany
Foliage Western Tent Caterpillars <i>Malacosoma californicum</i> 31
<i>Clastoptera</i> spp
Cupressus Arizonica—Arizona cypress
Branchea & Terminals
Cedar Bark Beetles <i>Phloeosinus</i> spp85
Dense Mistletoe <i>Phoradendron densum</i> 184
Juniper Twig Pruner Styloxus Bicolor
Stem
Bark Beetles & Wood Borers
Juniper Borers <i>Trachykele blondeli</i> , etc 90
Rusts
Gymnosporangium Rusts Gymnosporangium spp 169
Fraxinus spp.—ash
Branches & Terminals
Big Leaf Mistletoe <i>Phoradendron macrophyllum</i> 184
Foliage
Fall Webworm <i>Hyphantria cunea</i>
Juglans major—Arizona walnut
Branches & Terminals
Big Leaf Mistletoe <i>Phoradendron macrophyllum</i> 184
Juniperus osteosperma—Utah Juniper, J. deppeana—alligator
juniper, J. monosperma—one-seed juniper, J. scopulorum—
rocky mountain juniper
Branches & Terminals
Cedar Bark Beetles <i>Phloeosinus</i> spp85
Juniper Mistletoe <i>Phoradendron juniperinum</i> 184
Juniper Twig Pruner Styloxus Bicolor 107

Foliage	
Spider Mites Oligonychus spp	. 54
Spittlebugs Aphrophora spp., Clastoptera spp	. 53
Stem	
Bark Beetles & Wood Borers	
Juniper Borers Trachykele blondeli, etc	. 90
Rusts	
Gymnosporangium Rusts Gymnosporangium spp	169
Picea engelmannii—Engelmann spruce	
Branches & Terminals	
Western Spruce Dwarf	
Mistletoe Arceuthobium microcarpum	177
Foliage	
Cooley Spruce Gall Adelgid Adelges cooleyi	. 40
Frost	206
Nepytia janetae	. 27
Pine Needle Scale Chionaspis pinifoliae	. 49
Spruce Aphid <i>Elatobium abietinum</i>	. 42
Western Spruce Budworm Choristoneura occidentalis	8
Roots	
Armillaria or Shoestring Root Rot Armillaria spp	191
Schweinitzii Root and Butt Rot Phaeolus schweinitzii	198
Tomentosus Root Disease Inonotus tomentosus	196
Stem	
Bark Beetles & Wood Borers	
Ambrosia Beetles	. 92
Spruce Beetle <i>Dendroctonus rufipennis</i>	. 74
Decays	
Indian Paint Fungus Echinodontium tinctorium	138
Red Ring Rot <i>Phellinus pini</i>	
Rusts	
Spruce Broom Rust Chrysomyra arctostanhyli	171

Picea pungens—Colorado blue spruce
Branches & Terminals
Western Spruce Dwarf
Mistletoe Arceuthobium microcarpum 17'
Foliage
Cooley Spruce Gall Adelgid Adelges cooleyi 40
Douglas-Fir Tussock Moth Orgyia pseudotsugae 1
Frost
Pine Needle Scale Chionaspis pinifoliae
Spruce Aphid <i>Elatobium abietinum</i> 42
Western Spruce Budworm Choristoneura occidentalis 8
Roots
Armillaria or Shoestring Root Rot Armillaria spp 19
Schweinitzii Root and Butt Rot Phaeolus schweinitzi 198
Tomentosus Root Disease <i>Inonotus tomentosus</i> 196
Stem
Bark Beetles & Wood Borers
Ambrosia Beetles92
Spruce Beetle <i>Dendroctonus rufipennis</i>
Decays
Red Ring Rot <i>Phellinus pini</i>
Rusts
Spruce Broom Rust <i>Chrysomyxa arctostaphyli</i> 17
Pinus aristata—bristlecone pine
Branches & Terminals
Western Spruce Dwarf
Mistletoe Arceuthobium microcarpum17
Pinus edulis—piñon, P. monophylla—singleleaf piñon, P.
cembroides—Mexican piñon
Branches & Terminals
Bark Moths Dioryctria spp
Pinyon Dwarf Mistletoe Arceuthobium divaricatum 17'
Pinyon Pitch Nodule Moth <i>Petrova arizonensis</i> 105

Pitch M	oth Synanthedon spp	106
Twig Be	eetles Pityophthorus spp., Pityogenes spp.,	
	rus secundus	83
Foliage		
Drough	t	205
	erma Needle Cast Elytroderma deformans	
•	<i>Neodiprion</i> Spp., <i>Zadiprion</i> spp	22
Pine-Fe	eding Needleminers Coleotechnites spp	14
	Needle Scale Matsucoccus acalyptus	
	Spindle Gall Midge <i>Pinyinia edulicola</i>	
•	nd Needle Blight <i>Dothistroma septospora</i>	
	Mites Oligonychus spp	
_	oth Lophocampa ingens	
Roots		
Black S	tain Root Disease Leptographium wageneri	200
Stem		
Bark Be	eetles & Wood Borers	
Pin	e Engravers <i>Ips</i> spp	59
	od Borers	
Decays		
Red Ro	t Dichomitus squalens	136
Pinus eldarica-	—Mondel pine	
Stem	•	
Rusts		
Cor	mandra Blister Rust Cronartium comandrae	167
Pinus engelma	nnii—Apache pine	
Branches &	Terminals	
Southw	estern Dwarf Mistletoe Arceuthobium	
vaginati	um subsp. cryptopodum	177
Stem	- ** *	
	eetles & Wood Borers	
Sou	thern Pine Beetle Dendroctonus frontalis	70

Pinus nexuis—ilmoer pine	
Branches & Terminals	
Limber Pine Dwarf Mistletoe	
Arceuthobium cyanocarpum17	17
Stem	
Bark Beetles & Wood Borers	
Mountain Pine Beetle Dendroctonus ponderosae 6	55
Red Turpentine Beetle <i>Dendroctonus valens</i>	12
Cankers	
Atropellis Canker Atropellis piniphila	58
Pinus leiophylla var. chihuahuana—Chihuahua pine	
Branches & Terminals	
Chihuahua Pine Dwarf Mistletoe Arceuthobium gillii 17	17
Cones & Seeds	
Southwestern Cone Rust Cronartium conigenum 17	15
Stem	
Bark Beetles & Wood Borers	
Southern Pine Beetle <i>Dendroctonus frontalis</i>	70
Pinus ponderosa var. arizonica—Arizona pine	
Branches & Terminals	
Southwestern Dwarf Mistletoe Arceuthobium	
vaginatum subsp. cryptopodum	17
Pinus ponderosa—ponderosa pine	
Branches & Terminals	
Bark Moths <i>Dioryctria</i> spp)6
Pine Tip Moths)1
Pitch Moth Synanthedon spp)6
Southwestern Dwarf Mistletoe Arceuthobium	
vaginatum subsp. cryptopodum	17
Twig Beetles Pityophthorus spp., Pityogenes spp.,	
Pityoborus secundus8	33
Western Pineshoot Borer <i>Eucosma sonomana</i> 10)3

Cones & Seeds	
Cone Beetles Conophthorus spp	95
Pine Coneworm Dioryctria auranticella	99
Pine Seed Chalcid Megastigmus albifrons	98
Ponderosa Pine Seedworm Cydia piperana	97
Foliage	
Aphids	38
Drought	05
Elytroderma Needle Cast <i>Elytroderma deformans</i> 12	23
Lophodermella Needle Cast Lophodermella cerina 12	
Pandora Moth Coloradia pandora	
Neophasia menapia	20
Pine Needle Scale Chionaspis pinifoliae	49
Neodiprion spp., Zadiprion spp	
Pine-Feeding Needleminers Coleotechnites spp	
Ponderosa Pine Resin Midge Cecidomyia piniinopis	
Prescott Scale Matsucoccus vexillorum	46
Red Band Needle Blight Dothistroma septospora 12	25
Red Belt	08
Salt Damage	11
Spider Mites Oligonychus spp	54
Tiger Moth Lophocampa ingens	24
Winter Injury	07
Roots	
Annosus Root Rot Heterobasidion annosum	93
Armillaria or Shoestring Root Rot Armillaria spp 19	91
Schweinitzii Root and Butt Rot Phaeolus schweinitzii 19	98
Stem	
Bark Beetles & Wood Borers	
Ambrosia Beetles	92
Mexican Pine Beetle Dendroctonus mexicanus	70
Mountain Pine Beetle Dendroctonus ponderosae 6	65
Pine Engravers <i>Ips</i> spp	59
Red Turpentine Beetle Dendroctonus valens	72
Roundheaded Pine Reetle Dendroctorus adjunctus	67

Western Pine Beetle <i>Dendroctonus brevicomis</i> 63	3
Wood Borers	3
Cankers	
Atropellis Canker Atropellis piniphila 158	3
Decays	
Pouch Fungus Cryptoporus volvatus	2
Red Rot Dichomitus squalens136	5
Other	
Carpenter Ants Camponotus spp)
Termites	ĺ
Rusts	
Comandra Blister Rust Cronartium comandrae 167	
Limb Rust Cronartium arizonicum	1
Western Gall Rust Peridermium harknessii 166	5
Pinus strobiformis—southwestern white pine	
Branches & Terminals	
Apache Dwarf Mistletoe Arceuthobium apachecum 177	7
Foliage	
White Pine Needle Cast Lophodermella arcuata 129)
Roots	
Schweinitzii Root and Butt Rot Phaeolus schweinitzii 198	3
Stem	
Bark Beetles & Wood Borers	
Mountain Pine Beetle <i>Dendroctonus ponderosae</i> 65	5
Red Turpentine Beetle <i>Dendroctonus valens</i> 72	2
Cankers	
Atropellis Canker <i>Atropellis piniphila</i> 158	3
Rusts	
White Pine Blister Rust Cronartium ribicola 163	1
Platanus wrightii—Arizona sycamore	
Branches & Terminals	
Big Leaf Mistletoe <i>Phoradendron macrophyllum</i> 184	1
Foliage	
Sycamore Anthracnose <i>Apiognomonia veneta</i>	3

Fotyporus	anceps See Dicholintus squalens
Populus s	pp.—cottonwood
Brancl	nes & Terminals
Bi	g Leaf Mistletoe <i>Phoradendron macrophyllum</i> 184
Foliag	е
Ap	ohids
Bl	ack Leaf Spot Marssonina populi
Fa	ll Webworm <i>Hyphantria cunea</i> 29
Inl	x Spot Leaf Blight Ciborinia whetzelii
Me	elampsora Rust Melampsora spp116
Sh	epherd's Crook Venturia spp117
We	estern Tent Caterpillars Malacosoma californicum 31
Stem	
Ba	rk Beetles & Wood Borers
	Ambrosia Beetles92
Ca	nkers
	Cytospora Canker Cytospora chrysosperma 152
Populus ti	remuloides—aspen
Foliag	<u> </u>
Bl	ack Leaf Spot <i>Marssonina populi</i>
	ought
Fre	ost
Inl	Spot Leaf Blight Ciborinia whetzelii
	rge Aspen Tortrix Choristoneura conflictana33
Me	elampsora Rust <i>Melampsora</i> spp
	epherd's Crook <i>Venturia</i> spp
Sp	ider Mites <i>Oligonychus</i> spp54
_	estern Tent Caterpillars Malacosoma californicum 31
Roots	
Ga	noderma Root Rot <i>Ganoderma applanatum</i> 202
Stem	• •
Ba	rk Beetles & Wood Borers
	Ambrosia Beetles92
	Wood Borers

Calikers	
Black Canl	ker Ceratocystis fimbriata148
Cryptospha	neria Canker Cryptosphaeria lignyota 154
Cytospora	Canker <i>Cytospora chrysosperma</i> 152
Hypoxylon	Canker Hypoxylon mamatum 156
Sooty Bark	Canker Encoelia pruinosa
Decays	-
False Tinde	er Conk <i>Phellinus tremulae</i> 144
Red Belt F	ungus Fomitopsis pinicola 140
<i>Prosopis</i> spp.—mesqu	ite
Branches & Terminals	
Desert Mistletoe Phora	ndendron californicum184
<i>Prunus virginiana</i> var	s. demissa and
melancocarpa—comn	
Foliage	, and the second
9	Hyphantria cunea29
Pseudotsuga menziesii	
Branches & Termin	
Douglas-Fir Dy	varf Mistletoe <i>Arceuthobium douglasii</i> 177
Foliage	O
•	Gall Adelgid Adelges cooleyi40
• •	ssock Moth Orgyia pseudotsugae 11
_	ast <i>Melampsora</i> spp116
	ir Looper <i>Galenara consimilus</i> 26
	ale <i>Chionaspis pinifoliae</i> 49
	eedle Cast <i>Rhabdocline</i> spp 126
	Budworm Choristoneura occidentalis 8
Roots	
Armillaria or S	hoestring Root Rot Armillaria spp 191
	ot Disease <i>Leptographium wageneri</i> 200
	oot and Butt Rot <i>Phaeolus schweinitzii</i> 198
Tomentosus Ro	oot Disease <i>Inonotus tomentosus</i> 196

Stem	
Bark Beetles & Wood Borers	
Ambrosia Beetles9	2
Douglas-Fir Beetle Dendroctonus pseudotsugae 7	7
Decays	
Indian Paint Fungus <i>Echinodontium tinctorium</i> 13	8
Pouch Fungus Cryptoporus volvatus	
Red Belt Fungus Fomitopsis pinicola	
Red Ring Rot <i>Phellinus pini</i>	
Other	
Carpenter Ants Camponotus spp 10	9
Termites	1
Quercus hypoleucoides—silverleaf oak	
Cones & Seeds	
Southwestern Cone Rust Cronartium conigenum 17	5
Quercus spp.—oak	
Branches & Terminals	
Oak Mistletoe <i>Phoradendron coryae</i> 18	4
Foliage	
Frost	6
Spider Mites Oligonychus spp5	
Spittlebugs <i>Aphrophora</i> spp., <i>Clastoptera</i> spp 5	
Roots	
Armillaria or Shoestring Root Rot Armillaria spp 19	1
Stem	
Decays	
Canker Rot of Oak <i>Inonotus andersonii</i>	6
Heart Rots of Oak <i>Inonotus dryophilus</i>	5
Heart Rots of Oak <i>Phellinus everhartii</i>	
Ribes spp.—currants and gooseberries	
Foliage	
White Pine Blister Rust Cronartium ribicola16	1

Salix spp.—willow	
Foliage	
Fall Webworm Hyphantria cunea	29
Melampsora Rust Melampsora spp	116
Western Tent Caterpillars Malacosoma californica	<i>ım</i> 31
Stem	
Cankers	
Cytospora Canker Cytospora chrysosperma	152

General Index

A	
Abiotic disorders	2
Adelgedae	8
Adelges cooleyi	0
Adelgid	
Cooley Spruce Gall4	0
Alder Flea Beetle	
Altica ambiens	5
Animal Damage	
Annosus Root Rot	
Aphid	
Conifers	8
Hardwoods	8
Spruce	2
Aphididae3	
Apiognomonia veneta	
Arceuthobium spp	
abietinum f. sp. concoloris	
apachecum	
blumeri	
divaricatum18	
douglasii	
gillii	
microcarpu	
vaginatum subsp. cryptopodum	
Armillaria or Shoestring Root Rot	
<i>Armillaria</i> spp	
Artist's Conk	
Atropellis piniphila	
_	
В	
Beetles, Ambrosia	
Beetles, Bark 55-8	
Cedar Bark Beetles	
7	
	8

Cerambydidae	88
Ceratocystis fimbriata1	48
Ceratocystis spp1	43
Chionaspis pinifoliae	49
Choristoneura	
conflictana	33
occidentalis	8
Chrysomyxa arctostaphyli1	71
Ciborinia whetzelii	15
Clastoptera	53
Coleotechnites spp	14
Coloradia pandora	18
Cone Insects	99
Coneworm, Pine	99
Conophthorus spp	95
Cow Patty. See Phaeolus schweinitzii	
Cronartium	
arizonicum	64
comandrae1	67
conigenum1	75
ribicola	61
Cryptoporus volvatus	42
Cryptosphaeria lignyota	54
Cydia piperana	97
Cytospora Canker	
Cytospora chrysosperma	52
D	
Decay	
Root Disease	03
Annosus	93
Armillaria	91
Ganoderma	02
Schweinitzii	98
Tomentosus 1	96

Stem Decays
False Tinder Conk144
Indian Paint Fungus
Oak
Pouch Fungus
Red Belt Fungus
Red Ring Rot
Red Rot
Defoliators
Dendroctonus
adjunctus67
brevicomis63
frontalis70
mexicanus70
ponderosae
pseudotsugae77
rufipennis
valens72
Dichomitus squalens
Dioryctria auranticella
<i>Dioryctria</i> spp
Dothistroma septospora
Drought
Dryocoetes confusus
, , , , , , , , , , , , , , , , , , ,
E
Echinodontium tinctorium
Elatobium abietinum
Elk
Elytroderma deformans
Encoelia pruinosa
Epinotia meritana
Fucosma sonomana 103

Flatheaded borers	88
Foliar Diseases	131
Fomitopsis pinicola	140
Frost	206
G	
Galenara consimilus	26
Gall Formers	37
Ganoderma applanatum	202
Ganoderma Root Rot	
Gymnosporangium spp	
н	
Hemiptera	37
Heterobasidion annosum	
Homoptera	
Hyphantria cunea	
Hypoxylon mamatum	
- Inonotus	
andersonii	146
dryophilus	
tomentosus	
Invasive pests	
<i>Ips</i> spp.	
confusus	
lecontei	
pini	
Isoptera	
wopieru	. 1 1
J	
Juniner Twig Pruner	107

K
Key to insects and diseases
L
Large Aspen Tortrix
Leaf Disease
Black Leaf Spot
Ink Spot Leaf Blight115
Melampsora Rust
Shepherd's Crook
Sycamore Anthracnose
Leptographium wageneri
Lightning210
Lirula abietis-concoloris
Looper
New Mexico Fir
Lophocampa ingens
Lophodermella arcuata129
Lophodermella cerina121
M
Malacosoma californicum
Marssonina populi
Matsucoccus
acalyptus44
vexillorum
Megastigmus albifrons
Melampsora spp
Melampsorella caryophyllacearum
Midge
Pinyon Spindle Gall
Ponderosa Pine Resin
Mistletoes
Dwarf Mistletoes
Apache dwarf mistletoe
Rlumer's dwarf mistletoe 181

		Chihuahua pine dwarf mistletoe
		Pinyon dwarf mistletoe
		Southwestern dwarf mistletoe
		Western spruce mistletoe
	True Mi	istletoes
		Bigleaf mistletoe
		Dense mistletoe
		Desert mistletoe
		Hairy juniper mistletoe
		Juniper mistletoe
		Southwestern oak mistletoe
		Texas juniper mistletoe
Mites		54
Moth		
Wioth	Bark	
	Dark	
	Pandora	ı,
		D
		Pitch Nodule
	Pitch	
	Tiger	
N		
Needle	Cost	
Needle		erma
		ermella
	Rea Bai	nd Needle Blight (Pine)
	т г.	120
NT 11		Fine
Needle		
	Pine-fee	eding
		9 10

Neodiprion spp22
Neophasia menapia20
Nepytia janetae
New Mexico Fir Looper
Non-infectious disorders
0
Oligonychus spp54
<i>Ophiostoma</i> spp
Orgyia pseudotsugae
P
Pandora moth
Peridermium harknessii
Petrova arizonensis105
Phaeolus schweinitzii
Phellinus
everhartii
pini
tremulae
Phloeosinus spp85
Phoradendron spp
californicum
capitellatum
coryae
densum
hawksworthii188
juniperinum188
<i>macrophyllum</i>
pauciflorum188
Pinyinia edulicola 52
Pityoborus secundus
Pityogenes spp
Pityophthorus spp
Porcupine

R		
Red Belt		309
Rhabdoo	<i>line</i> spp	26
Rhyacio	<i>nia</i> spp	01
Root Dis	ease	203
Roundhe	aded borers	88
Rust		75
(Comandra Blister Rust	
I	ir Broom Rust	71
(Symnosporangium Rusts	69
I	imb Rust	64
5	outhwestern Cone Rust	75
	pruce Broom Rust	
	Vestern Gall Rust	
7	Vhite Pine Blister Rust	60
S		
	nage2	11
Sapsuck	e	
•	king Insects	
Sup Suc	ang mocets 37	
F	ine	22
Scale		
I	ine Needle	49
F	inyon Needle	44
F	rescott	46
Schwein	itzii Root and Butt Rot	98
Scolytus	ventralis	81
Seed Ch	alcid, Pine	98
Seed Ins	ects	.99
Seedwor	m	
I	Onderosa Pine	97
Shoot In	sects	07
Spider M	lites	54
Snittlehr	σς	53

Stain										
Black		 	 	 	 	 				. 200
Stains		 	 	 	 	 	 			84
Stem Decays		 	 	 	 	 	 		132	2-146
Styloxus bicolo	r	 	 	 	 	 	 		. 53	, 107
Synanthedon sp	р	 	 	 	 	 				. 106
т										
Termites		 	 	 	 	 	 			. 111
Tiger moth										
Tomentosus Ro										
U										
Ungulates		 	 	 	 	 	 			. 213
V										
Venturia spp.		 	 	 	 	 				. 117
W										
Webworm										
Fall		 	 	 	 	 	 			29
Western Spruce										
Western Tent C										
White Pine Blis										
Winter Injury										
wood decay		 	 	 	 	 	 		133	, 191
Z										
Zadiprion spp										22